FSN-FG 0001

SURVIVAL, GROWTH AND FEED UTILIZATION OF THE RECIPROCAL HYBRIDS OF *Clarias gariepinus* (BURCHELL, 1822) AND *Heterobranchus bidorsalis* (GEOFFROY, 1809) IN CONCRETE TANKS

OWODEINDE, F.G.¹, NDIMELE, P.E.^{1*}, JENYO-ONI, A.² and ONYENANIA, O.B.¹

¹Department of Fisheries, Faculty of Science, Lagos State University, Ojo, Lagos, Nigeria.

²Department of Wildlife and Fisheries Management, Faculty of Agriculture and Forestry, University of Ibadan

Copyright 2010, Fisheries Society of Nigeria.

This paper was prepared for presentation at the 25th Annual International Conference and Exhibition in Administrative Staff College of Nigeria (ASCON), Topo-Badagry, Lagos, Nigeria, 25th – 29th October, 2010.

This paper was selected for presentation by an FISON Program Committee following review of information contained in an abstract submitted by the author(s). Contents of the paper, as presented, have not been reviewed by the Fisheries Society of Nigeria and are subject to correction by the author(s). The material, as presented, does not necessarily reflect any position of the Fisheries Society of Nigeria, its officers, or members. Papers presented at FISON meetings are subject to publication review by Editorial Committees of the Fisheries Society of Nigeria. Electronic reproduction, distribution, or storage of any part of this paper for commercial purposes without the written consent of the Fisheries Society of Nigeria is prohibited. Permission to reproduce in print is restricted to an abstract of not more than 300 words; illustrations may not be copied. The abstract must contain conspicuous acknowledgement of where and by whom the paper was presented. Write Librarian, Fisheries Society of Nigeria (FISON), P. O. Box 2607 Apapa, Lagos.

Abstract

*An experiment was conducted to determine the survival rate, fertilization rate, growth performance and feed utilization of the reciprocal hybrids of *Clarias gariepinus* and *Heterobranchus bidorsalis*. Two genetic crosses were made: *C. gariepinus*♀ x *H. bidorsalis*♂ (*clariabanchus*) and *H. bidorsalis*♀ x *C. gariepinus*♂ (*heteroclarias*). The experiment was divided into two phases; artificial propagation of the fish species using synthetic hormone and rearing the fry for 14 days; and rearing the 14 days old fry for 35 days. In the first phase of the experiment, survival of fry were estimated in each experimental unit (genetic cross) while in the second phase, growth and nutrient utilization were investigated. The result revealed that the highest %fertilization, hatching rate and %survival occurred in *H. bidorsalis*♀ x *C. gariepinus*♂ (*heteroclarias*) and the differences were significant ($p < 0.05$). Percentage weight gain and specific growth rate were significantly ($p < 0.05$) higher in *clariabanchus* than *heteroclarias*. Based on the result of this study, reciprocal hybrids of *Clarias**

gariepinus and *Heterobranchus bidorsalis* is recommended for commercial aquacultural practices.

Keywords: African catfish, survival rate, growth performance, feed utilization, reciprocal hybrid

Introduction

Fish is an important and cheapest source of animal protein it accounts for about 37% of Nigeria's total protein requirement {Federal Department of Fisheries (FDF), 2002}. Fish production in Nigeria is mainly from the captured sector, especially artisanal coastal and artisanal inland fisheries. This sector contributes over 80% of total domestic production of about 510,000 tonnes per annum (FDF, 2004). Over-exploitation of the marine fishery resources has resulted in gradual depletion of the stock. FAO (2003) reported that Nigeria is one of the largest importers of fish in the developing world, importing about 600,000 metric tonnes annually. To solve this short-fall in fish supply, Nigeria must encourage investments in aquaculture.

Genetic technologies have been used in different areas of biology to develop species that possess desirable characteristics for culture. Bartley *et al* (1997) reported that the hybridization of the African catfish (*Clarias gariepinus*) with the Thai catfish (*Clarias macrocephalus*) produced offspring that was preferred to the parental species because they had the desired flesh quality of the Thai catfish and the fast growth of the African catfish. Legendre *et al* (1992) reported that *Heterobranchus* species had

a growth rate twice as fast as that of *Clarias gariepinus* while *Clarias gariepinus* survives in poorly oxygenated water (Teugels, 1996). Hybridization between these two clariid catfishes may produce offspring that possess a combination of these desirable qualities.

The continuous growth of aquaculture is hinged on the production of fish seeds with high fertilization and survival rates, high feed conversion efficiency, and high growth rate among other factors (Adebayo and Popoola, 2008). This kind of seed can be obtained through genetic technology. This study investigates the fertilization, hatching and survival rates of larvae of reciprocal hybrids of two African clariid catfish (*Clarias gariepinus* and *Heterobranchus bidorsalis*) and also studies their growth performance and feed utilization.

Materials and Methods:

Experimental Fish:

This study was conducted in 2007. Sexually matured brood fish (500 - 600g) were obtained from Lagos State University, Nigeria. The brood fish consist of two male each of *Clarias gariepinus* and *Heterobranchus bidorsalis* and two female each of this same fish species. The broodstocks were selected based on their external morphological features as described by Viveen *et al* (1985).

Experimental Procedure:

The experiment was divided into two broad stages;

- (a) Artificial propagation of fish species, using synthetic hormone and raising the fry for 14 days.
- (b) Rearing of 14days old fry for another 35days.

(A) Artificial propagation

The broodstocks used for the experiment were conditioned for two weeks in holding tanks in the hatchery of Lagos State University and were fed 40% crude protein pelleted feed at 3% body weight twice daily at 09:00 hr and 16:00 hr.

The female broodstocks were induced by injecting Ovaprim, a synthetic hormone (Aqualife Syndel International Inc. Vancouver, B.C. Canada) at the rate of 0.5ml / 1000g body weight. Ovulation occurred -15 – 18 hrs after injection and gentle pressure was applied to the anterior-posterior direction on the abdomen of the female brood fish (*C. gariepinus* and *H. bidorsalis*) to strip them of eggs. The male brood fish of (*C. gariepinus* and *H. bidorsalis*) were anaesthetized, sacrificed and their testes removed. Milt was collected after dissection of the testes and immediately preserved in 0.9% NaCl solution. Stripped eggs were later fertilized with milt after sperm activation was initiated by the addition of 5ml fresh water and checked for motility by microscopic examination (Viveen *et al*, 1985). These are the two (2) crosses:

- (1) *Clarias gariepinus*♀ x *Heterobranchus bidorsalis*♂ (Clariabanchus)
- (2) *Heterobranchus bidorsalis*♀ x *Clarias gariepinus*♂ (Heteroclarias)

After about 1min of gentle stirring, fertilized eggs were rinsed in fresh water to remove excess milt and treated with talcum powder for 15-30 minutes to inhibit adhesiveness of the egg jelly coat and to prevent clumping and suffocation of eggs during incubation. Eggs were incubated in glass tank (70 x 45 x 35 cm³). The incubation jars was aerated continually and temperature was 27±1°C. Hatching occurred 23-26 hrs later. The larvae were left for three days in the incubation jars to absorb their yolk.

After yolk absorption, the post-larvae were fed *Artemia* naupli for a period of 14days. Aeration was done continually and the water temperature, pH and dissolved oxygen were 27±1°C, 6.5 and 4.5 mg/l respectively. Water was changed regularly to avoid mortality resulting from polluted water.

(B) Growth Experiment

A total of 180 14days old hybrid catfish (Heteroclarias and Clariabanchus) juveniles were used. Thirty specimens of each hybrid were randomly chosen and

allocated to 6 circular flow-through tanks {each experimental unit (that is genetic cross) had three replicates}. Rearing conditions were the same as described above. The diameter of each tank was 2m and there was at least 50% water exchange daily. Each tank contained about 45l of fresh water.

Prior to stocking, quicklime was applied to the tank bottom at 150 g/m² to eliminate parasites and invertebrate predators.

Feeding trials:

The fish in each of the culture treatments was fed on commercial pelleted diet (56% crude protein) (Table 1) at 3% of their body weight according to the recommendation of Viveen *et al* (1985). The daily ration was divided into two; one part was fed at 09:00 hr and the other part at 18:00 hr. Feed were dispensed evenly on the water surface in each tank to allow equal opportunity for feeding. Feeding in all tanks was generally completed in about 10-15 minutes. Weighing of fish was carried out weekly throughout the period of the experiment. On weighing days, fish were not fed until the whole exercise was completed. Feeding rate was recalculated to accommodate for the weight changes. The feeding trials lasted for five weeks.

Determination of water quality parameter:

Temperature of water in all tanks was measured daily using mercury-in-glass thermometer. The pH was measured by a pH meter (Jenway model 9060). Dissolved oxygen concentration in water was determined weekly using the methods of APHA (1989).

Reproductive performance parameter:

The number of eggs released in each treatment unit was determined by subtracting the weight of the brood stock after stripping (W_b) in grams from the weight of the brood stock before stripping (W_a) in grams and multiplying the difference by 700 (1gm = 700 eggs) (Viveen *et al*, 1985). Fertilization rate was determined when eggs generally reached the 4-8 celled stage of embryonic

development. For calculating percentage fertilization, a sample of about 50eggs from each replicate of each treatment were carefully taken on Petri dish containing water and counted under a microscope (40 times magnification) (Adebayo, 2006). The fertilization rate was then calculated based on the total number of eggs counted. After hatching, the numbers of larvae within each experimental unit were carefully counted and the hatching rate was calculated. Similarly, the survival rate was calculated at the end of the rearing period (49 days after hatching) based on the initial number of larvae used in the experiment.

Growth and Nutrient Utilization:

Growth and nutrient utilization were analyzed by calculating the weight gain (WG) over the test period, percent weight gain (PWG), specific growth rate (SGR), food conversion ratio (FCR) and protein efficiency ratio (PER).

$$\text{Specific growth rate (SGR)} = \frac{\text{Log}_e W_2 - \text{Log}_e W_1}{T_2 - T_1} \times \frac{100}{1}$$

Where W_2 = Weight of fish at time T_2 in days; W_1 = Weight of fish at time T_1 in days; Log_e = Natural Logarithm.

Food conversion ratio (FCR) as determined by Weight of dry feed fed (g) divided by live weight gain (g); and

Protein efficiency ratio (PER) defined as gain in weight of test fish (g) divided by the amount of Protein consumed (g).

Statistical Analysis:

Statistical analysis was performed using the SPSS V. 15.0 package for Windows. Analysis of Variance (ANOVA) was used and where significant difference was indicated, means were tested using Fishers Least Significant Difference (LSD) test at $p = 0.05$ significance level.

Result and Discussion

The fertilization, hatching and survival rates of catfish hybrids (*Clariabanchus* and *Heteroclarias*) investigated in this study are generally high (Table 2) except the hatching rate of *C. gariepinus*♀ x *H. bidorsalis*♂ (*Clariabanchus*) (52.57 ± 0.88). This may be due to the tolerable physico-chemical qualities of the culture water (Aliu and Obasogie, 2006). The hybrid (*H. bidorsalis*♀ x *C. gariepinus*♂, *Heteroclarias*) had the best fertilization, hatching and survival rates.

Although, the fertilization, hatching and survival rates of *H. bidorsalis*♀ x *C. gariepinus*♂ (*heteroclarias*) was significantly ($p < 0.05$) higher than *C. gariepinus*♀ x *H. bidorsalis*♂ (*clariabanchus*), the values obtained in this study are lower than those reported in previous studies (Adebayo, 2006; Ataguba *et al.*, 2009). The relatively lower fertilization, hatching and survival rates recorded in this study might be due to differences arising from breeding history, age, water quality and season (De Graaf *et al.*, 1995 and Ataguba *et al.*, 2009). Percentage weight gain and specific growth rate of *clariabanchus* were significantly ($p < 0.05$) higher than that of *heteroclarias* (Table 3). However, these results could not be confirmed with the result of the mean weight gain (MWG), feed conversion ratio (FCR) and protein efficiency ratio (PER) because the difference in these parameters were not significant ($p > 0.05$).

CONCLUSION

This study has confirmed earlier studies that cross breeding of African clariid catfish can be done successfully and the products of the cross breed can also be reared to adults.

ACKNOWLEDGEMENT

The authors are grateful to the Head, Department of Fisheries, Faculty of Science, Lagos State University, Ojo, Lagos, Nigeria for approving the use of the

Laboratory and providing the necessary materials for the study.

REFERENCES

- Adebayo, O.T., (2006): Reproductive performance of African Clariid Catfish *Clarias gariepinus* broodstock on varying maternal stress. *Journal of Fisheries International* 1(1- 2), 17-20.
- Adebayo, O.T.; Popoola, O.M., (2008): Comparative evaluation of efficacy and cost of synthetic and non-synthetic hormones for artificial breeding of African catfish (*Clarias gariepinus* Burchell, 1822). *Journal of Fisheries and Aquatic Sciences*, 3(1), 66-71.
- Aliu, B.S.; Obasogie, B.E., (2006): The growth responses of African catfish, *Heterobranchus* x *Clarias* hybrid fingerlings fed dietary plant and animal lipids. *Nigerian Journal of Fisheries*, 2/3 (2), 312-319.
- APHA, American Public Health Association (1989): American Water Works Association and Water Pollution Control Federation (1989) Standard method for the examination of water waste water (17th edition), APHA, New York U.S.A. 1268 pp,
- Ataguba, G.A., Annune, P.A.; Ogbe, F.G., (2009): Induced breeding and early growth of progeny from crosses between two African clariid fishes, *Clarias gariepinus* (Burchell) and *Heterobranchus longifilis* under hatchery conditions. *Journal of Applied Biosciences* 14, 755-760.
- Bartley, D.M.; Rana, K. and Immink, A.J., (1997): The use of inter-specific hybrids in aquaculture. The FAO Aquaculture Newsletter No 17, Rome, Italy. 7-13. December, 1997. ISSN 1020-3443.
- De Graaf, G.J.; Galemoni, F.; Banzoussi, B., (1995): Artificial reproduction and fingerling production of the

- African Catfish, *Clarias gariepinus* (Burchell 1822) in protected and unprotected ponds. *Aquaculture Research* 26, 233-242.
- Federal Department of Fisheries (FDF) (2002): Fisheries statistics of Nigeria.
- Federal Department of Fisheries (FDF) (2004): Fisheries statistics of Nigeria.
- Food and Agriculture Organisation (FAO) (2003): Opportunities for aquaculture development in Nigeria. FAO Aquaculture News letter No 17: 28pp.
- Legendre, M.; Teugel, G.G.; Carty, G.; Jalabart, B., (1992): A comparative study on morphology, growth rate and reproduction of *Clarias gariepinus* (Burchell 1822), *Heterobranchus longifilis* (Valenciennes 1840) and their reciprocal hybrids (Pisces: Clariidae). *Journal of Fish Biology* 40, 59-79.
- Teugels, G.G., (1996): Taxonomy, phylogeny and biogeography of catfishes (Ostariophysi; Siluroidei): an overview. *Aquatic Living Resources*. 9, 9-34.
- Viveen, W.J., Richter, C.J. Van Oodot, P.G. Janssen, J.A.; Huisman, E.A., (1985): Practical manual for the culture of the African Catfish, *Clarias gariepinus*. Section for Research and Technology, Ministry for Development Co-operation. The Hague, Netherlands. 128pp.

Table 1: Nutrient composition of commercial feed (CATCO FISH CONCENTRATE – COPPENS) fed to fry of reciprocal hybrids of clariid catfish.

Nutrient	% Composition
Crude protein	56.0
Crude fibre	10.9
Crude fat	15.0
Ash	10.9
Phosphorus	8.0
Energy	3400Kcal/Kg

*Each kg of the diet contains 300mg vit C, 200mg vit E, 22,500 IU vit A, 2,500 IU vit D₃, 5mg Cu, E280 Preservatives and E324 Anti-oxidants.

Table 2: Percentage fertilization, hatching rate and survival rate (14 days post hatching) of reciprocal hybrids (Clariabbranchus and Heteroclarias) of *Clarias gariepinus* and *Heterobranchus bidorsalis*

❖ Values in the same column and with the same superscript are not significantly different (p>0.05)

	% Fertilization (mean ± se)	Hatching rate (mean ± se)	%Survival (mean ± se)
<i>C. g</i> ♀ x <i>H. b</i> ♂ (n = 175)	65.87± 2.49 ^a	52.57± 0.88 ^a	65.22± 1.30 ^a
<i>H. b</i> ♀ x <i>C. g</i> ♂ (n = 350)	79.44± 5.39 ^b	70.57± 1.04 ^b	76.11± 1.59 ^b

Table 3: Growth of reciprocal hybrids of *Clarias gariepinus* and *Heterobranchus bidorsalis* between 14th and 49th day after hatching.
SGR = Specific Growth Rate; FCR = Food Conversion Ratio; PER = Protein Efficiency

Ratio.

➤ Values in the same column and with the same superscript are not significantly different (p>0.05)

Parameter	<i>C. g</i> ♀ x <i>H. b</i> ♂ (Clariabanchus)	<i>H. b</i> ♀ x <i>C. g</i> ♂ (Heteroclarias)
Initial Weight	0.52 ± 0.07	0.67 ± 0.11
Mean Weight.	1.31 ± 0.06 ^a	1.19 ± 0.05 ^a
Gain (g)		
Average Daily	0.04 ± 0.01 ^a	0.03 ± 0.01 ^a
Weight Gain (g)		
% Weight	2525.00 ± 109.67 ^a	1362.07 ± 101.95 ^b
Gain (%)		
SGR	9.34 ± 0.77 ^a	7.66 ± 0.79 ^b
% day ⁻¹)		
FCR	0.23 ± 0.18 ^a	0.35 ± 0.26 ^a
PER	11.92 ± 4.76 ^a	9.69 ± 3.45 ^a