Mesh-size selectivity of multifilament gill net at Fakun village, north of Lake Jebba

Ago, N. D. / Kwen, K. / Binyotubo, T. E.

Abstract

Studies on the mesh size selectivity of multifilament gillnet at Fakun village North of Lake Jebba were carried out. Four different mesh sizes (76mm, 101mm, 126mm and 177mm) of multifilament nets were used in the construction of the gillnet with hanging ratio 0.5 each. Daily catch from the nets were examined for seven weeks. The fish caught comprised of nine (9) species belonging to six (6) families. The result of analysis of variance of the number of species caught showed significant difference ($P < 0.05$) among the different mesh sizes of the gillnet. The catch mean weight for each mesh size (76mm, 101mm, 126mm and 177mm) was 266.91, 285.54, 305.10 and 349.02 respectively also the relative percentage of number of species caught by each of mesh size was 39.48% for 76mm, 23.58% for 101mm, 22.44% for 126mm and 14.48% for 177mm, which depicts the selectivity in the catch of the gillnets in this experiment.

Keywords: Mesh size selectivity, gillnet, Fakun village, Lake Jebba.

Introduction

The effectiveness of a gillnet depends on various factors which include mesh size, exposed net area, location, mesh shape, hanging ratio, visibility and type of netting materials in relation to stiffness and breaking strength (Van Brandt, 1984). Gillnet is very popular among artisanal fishermen in Nigeria. More than 75% of the fishermen in the inland coastal water in Nigeria uses gillnet at one time or another within a fishing season (Reed et al. 1967).

Knowledge of the efficiency of gillnet is important for estimation of fish population in stock assessment. The effects of technical innovation by fishers on the efficiency of gillnet is quantified for proper fisheries management. Netting material type has been shown to greatly influence catches. Transparent monofilament netting material is effective as gillnet in clear water. This is because the net is invisible to the fish. Visible nets tend to be avoided by fish. In turbid water however, the difference in structure and colour of gillnet material is usually very small (Nedelec, 1975).

Gillnets are passive gears, but can be used as active gears by dragging through water with the aid of two boats. They can be set passively on the surface, middle or bottom of water. The catchability and selectivity of gillnets depend on its hanging. One way of estimating the selectivity of gillnet is by comparing the catch with that of a relatively unselective gear such as the trawl net or the purse seine (Hamely 1975).

At Fakun fishing village, the people indulge in fishing practice at both subsistence and commercial level and in doing this they use different mesh size of gillnet ranging from less than 1" (25mm) and above which has the ability of catching young and juvenile fish. With this in mind, this study was carried out in order to:

1. Assess the size selection potential of multifilament gillnet using 3" (76mm) 4" (101mm) 5" (126mm) and 7" (177mm) mesh size.
2. Assess the efficiency of each of the mesh size based on their catch per unit effort.
3. Make necessary recommendation for the appropriate mesh size suitable for the water body.

Materials and Methods

Study area

The study was conducted at Fakun fishing village north of Jebba Lake, which is in the savannah zone, North-western Nigeria.

Construction Material and Fabrication Method

Four gillnets each measuring 15 metres in length and 3-metre depth were constructed using the following materials; white nylon (polyamide - PA) multifilament netting of twine-size 210d/2 (46tex) and mesh sizes of 76 mm, 101 mm, 126mm, and 177 mm and mounted at 50% ($E = 0.5$).

Line and ropes. The same sizes of head and foot-ropes were used for the four nets. This was made of *Kuralon* (polyvinyl-alcohol (PVA)) rope of thickness No. 8. **Accessories.** Cylindrically shaped synthetic corks of dimension 70 mm x 30 mm and each weighing 30g were used as float for all the net and were spaced at 5 metres interval. Flat lead sheet was caught into pieces of dimension 150 mm x 30 mm each weighing 100g and attached to the foot rope as sinkers with spacing interval corresponding to that of floats. **Experimental design.** The nets were ganged together randomly using randomized numbering technique to form a fleet of four gangs of net. The nets were set in the evening (between 1730 and 1830 GMT) and hauled in the following morning (between 0800 and 0900 GMT) thereby maintaining a soaked time of about 15hrs. **Data collection.** The catch from all nets were carefully removed as the nets were been hauled into the boat, separated according to the net's mesh size and sorted into species using fish identification keys prepared by Olaosebikan and Raji (2004) as taxonomic guide.

Results

Fish catch. Nine fish species belonging to six families were caught (Table 1). These are; *Brycinus nurse*, *Alestes brevis*, *Distichodus rostratus*, *D. brevipinnis*, *Citharinus citharus*, *Bagrus bayad*, *Synodontis membranaceus*, *Tilapia zilli* and *Oreochromis niloticus*. **Number and Biomass of fish caught.** The number and biomass of fish species caught are shown on Table 2. The total number of fish caught was 352 of which the highest relative percentage (39.48%) was recorded by net with 3" (76mm) mesh size followed by 4" (101mm) with 23.58% and 22.44% for 5" (126mm) mesh size. The least relative percentage of 14.48% was recorded from the net with 7" (177mm) mesh size. A total fish weight of 102.70kg was caught (Table 2). The highest relative percentage (36.12%) was recorded from the net with 3" (76mm) mesh size followed by 5" (126mm) mesh size with 23.47% and 4" (101mm) mesh size with 23.08%. The least relative percentage of total weight of fish caught was recorded from the net with 7" (177mm) mesh size (17.33%). The mean weights for 3" (76mm), 4" (101mm), 5" (126mm) and 7" (177mm) were 266.91g, 285.40g, 305.10g, and 349.02g respectively. The species diversity index were 1, 1, 0.88 and 0.77 for 3" (76mm) 4" (101mm) 5" (126mm) and 7" (177mm) respectively. Table 3 shows the relationship between the total number of fish caught and percentage mode of capture and it was observed that the higher relative percentage for mode of capture was recorded by entangling with 74.08%, while gilling had the least with 25.92%. The result of analysis of variance of the number of fish species caught by each mesh size showed there was significant difference ($P < 0.05$).

Discussion

The fish species caught in each of the mesh-size nets (Table 1) shows a diverse fishery nature. The difference in relative quantities of each type of fish caught by each mesh size of the gill nets could be attributed to the selectivity of the gill net due to differences in mesh size. Comparison of the result of the quantities of fish caught by each of the nets' mesh size (Table 2) show that the highest by number and biomass was recorded in the net with 3" (76 mm) mesh size. This might probably be due to the mesh size of the net which has the ability of catching only those species of fish that had grown up to it mesh size which were more compare to other fish sizes. The effectiveness of the gear to select a particular species had been found to be a product of two probabilities:

1. That of encountering of the gear by fish.
2. That of retaining of caught fish by the gear.

In this study, the population of fish in the Lake was unknown so the retention rate of the net was the factor used in estimating the catch characteristics of each mesh size of the gill nets. There is significant variation in the mean weight of fish caught in each net type. The 3" (76mm) gill net which caught more fish than 7" (177mm) gill net had a lower mean weight (266.91g) than the 7" (177mm) gill net (349.02). This indicates the ability of the larger mesh size net to retain large sized fish. Ita (1998) and du Feu and Abiodun (1999) gave a trend which show that the larger the mesh size of gill net, the bigger the size of fish caught.

The slight difference in species diversity index of the fish caught in 5" (126mm) and 7" (177mm) mesh size nets (0.88 and 0.77 respectively) depicts that both nets are comparable with relation to species selection properties.

Conclusion and Recommendation

In view of the result from this study, the gill net with mesh size 3" (76mm) is considered the best among others, because it was more efficient both in mode of capture and selectivity of different species of fish. It therefore recommended for artisanal fishers in Lake Jebba to improve their catch and leaving a balance ecosystem. However, there is need to modify the mesh size used to prevent the catching of undersized, juvenal fish.

Fig. 1 Map of Jebba Lake with Fakun at the extreme north.

Appendices

Table 1: Fish species caught by the gillnet

Family	Species
1. Characidae	<i>Brycinus nurse</i> , <i>Alestes brevis</i> .
2. Distichodontidae	<i>Distichodus rostratus</i> and <i>Distichodus brevipinnis</i>
3. Citharinidae	<i>Citharinus citharus</i>
4. Bagridae	<i>Bagrus bayad</i>
5. Mochokidae	<i>Synodontis membranaceus</i>
6. Cichlidae	<i>Tilapia zillii</i> , and <i>Oreochromis niloticus</i>

Table 2: Summary of the number and weight of fish species caught in all the mesh sizes during the experimental period.

Species	3" (76mm)		4" (126mm)		5" (177mm)		7" (177mm)		Overall total	
	No	Wt (g)	No	Wt (g)	No	Wt (g)	No	Wt (g)	No	Wt (g)
B. nurse	4	1600	2	1000					6	2600
A. brevis	5	2000	3	1200	1	600			9	3800
D. rostratus	10	6000	2	1500	1	500	1	800	14	8800
D. brevipinnis	8	4000	3	2000	3	2500	2	1000	16	9500
C. citharus	12	6000	6	3500	6	4000	3	3000	27	16500
B. bayad	3	3000	3	3500	4	4000	6	4000	16	14500
S. membranaceus	30	8000	25	7000	25	7500	17	7000	97	29500
T. zillii	32	3000	20	2000	21	2500	10	1000	83	8500
O. niloticus	35	35000	19	2000	18	2500	12	1000	84	9000
Total	139	37100	83	23700	79	24100	51	17800	352	102.70 kg
Mean weight		266.91		285.54		305.10		349.02		
Relative percentage	39.48	36.12	23.58	23.08	22.44	23.47	14.48	17.33		
No. of species	9		9		8		7			
Species diversity index	1		1		0.88		0.77			

Table 3: Relationship between total number of fish caught and percentage mode of capture

Species	Total	Gilling/Wedging		Entangling/Enmeshing	
	No.	No	%	No	%
<i>Brycinus nurse</i>	6	3	3.37	3	1.14
<i>Alestes brevis</i>	9	5	5.62	4	1.52
<i>Distichodus rostratus</i>	14	4	4.49	10	3.80
<i>Distichodus brevipinnis</i>	16	5	5.62	11	4.18
<i>Citharinus citharus</i>	27	7	7.87	20	7.60
<i>Bagrus bayad</i>	16	10	11.24	6	2.28
<i>Synodontis membranaceus</i>	97	50	56.18	47	17.87
<i>Tilapia zillii</i>	83	3	3.37	80	30.42
<i>Oreochromis niloticus</i>	84	2	2.23	82	31.18
Total	352	89		263	
Relative percentage			25.92		74.08

REFERENCES

- Du Feu, T. A. and Abiodun, J. (1999) Fisheries statistics of Kainji Lake, Northern Nigeria. Nov. 1994 to Dec. 1998. Nigeria–German (GTC) Kainji Lake fisheries Promotion Project. *Technical report Series No.13*. April. 128P.
- Hamley, J. M. (1975). Review of gillnet selectivity. *J. Fisheries Research Board Canada* 32: 1943–1969.
- Ita, E.O. (1998). The current status of fish stocks and fisheries in Kainji Lake, 1996 Nigeria–German. Kainji Lake Fisheries Promotion Project. *Technical Report Series*. December 1998. 45pp.
- Nedelec, C. (1975). FAO Catalogue of Small Scale Fishing Gear. Fishing News Book, London, England, by arrangement with Food and Agriculture Organization of the United Nations. 191pp.
- Olaosebikan, B. D. and Raji, A. (2004). Field Guide to Nigerian Freshwater Fishes–*Second Edition*. Federal College of Freshwater Fisheries Technology, New Bussa, Nigeria. 111pp.
- Reed, W., Hopson, J. B., Jonathan, A.J., Yaro, I. (1967). *Fish and Fisheries of Northern Nigeria* (1st Ed.). Ministry of Agriculture, Northern Nigeria. 222pp.
- Van-Brandt, A. (1984). Entangling nets. *Fish Catching Methods of the World* (Revised Ed.). London: Fishing New Books, 418P.