



UNIVERSIDADE ESTADUAL DE MARINGÁ
CENTRO DE CIÊNCIAS BIOLÓGICAS
DEPARTAMENTO DE BIOLOGIA
PROGRAMA DE PÓS-GRADUAÇÃO EM ECOLOGIA DE
AMBIENTES AQUÁTICOS CONTINENTAIS

LUCAS MACHADO NUNES

**Threatened, rare, and functionally rare species of fish in the
Upper Paraná River floodplain: ecological perspectives
and conservation**

Maringá
2023

LUCAS MACHADO NUNES

**Threatened, rare, and functionally rare species of fish in the
Upper Paraná River floodplain: ecological perspectives
and conservation**

Dissertação apresentada ao Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais do Departamento de Biologia, Centro de Ciências Biológicas da Universidade Estadual de Maringá, como requisito parcial para a obtenção do título de Mestre em Ecologia e Limnologia.

Área de Concentração: Ecologia e Limnologia

Orientador: Prof. Dr. Luiz Carlos Gomes

Maringá
2023

"Dados Internacionais de Catalogação na Publicação (CIP)"
(Biblioteca Setorial - UEM. Nupélia, Maringá, PR, Brasil)

N972t Nunes, Lucas Machado, 1988-
Threatened, rare, and functionally rare species of fish in the Upper Paraná River floodplain : ecological perspectives and conservation / Lucas Machado Nunes. -- Maringá, 2023.
41 f. : il. (algumas color.).
Dissertação (mestrado em Ecologia de Ambientes Aquáticos Continentais)-- Universidade Estadual de Maringá, Dep. de Biologia, 2023.
Orientador: Prof. Dr. Luiz Carlos Gomes.

1. Peixes de água doce - Espécies raras - Planície de inundação - Alto rio Paraná. 2. Peixes de água doce - Espécies ameaçadas de extinção - Planície de inundação - Alto rio Paraná. 3. Peixes de água doce - Comunidades, Ecologia de - Raridade taxonômica e funcional - Planície de inundação - Alto rio Paraná. 4. Biologia da conservação. I. Universidade Estadual de Maringá. Departamento de Biologia. Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais.

CDD 23. ed. -597.16809816

LUCAS MACHADO NUNES

**Threatened, rare, and functionally rare species of fish in the
Upper Paraná River floodplain: ecological perspectives
and conservation**

Dissertação apresentada ao Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais do Departamento de Biologia, Centro de Ciências Biológicas da Universidade Estadual de Maringá, como requisito parcial para a obtenção do título de Mestre em Ecologia e Limnologia e aprovada pela Banca Examinadora composta pelos membros:

BANCA EXAMINADORA

Prof. Dr. Luiz Carlos Gomes
Nupélia/Universidade Estadual de Maringá (Presidente)

Prof. Dr. Antonio Carlos Beaumord
Universidade do Vale do Itajaí (UniVali)

Prof. Dr. Matheus Tenório Baumgartner
Universidade Estadual de Maringá (UEM)

Prof.^a Dr.^a Elaine Antoniassi Luiz Kashiwaqui
Universidade Estadual do Mato Grosso do Sul (UEMS)

Dr. Fernando Miranda Lansac-Tôha
Nupélia/Universidade Estadual de Maringá (Pós-doutorando PEA/UEM)

Aprovada em: 18 de dezembro de 2023.

Local de defesa: Anfiteatro Prof. “Keshiyu Nakatani”, Nupélia, Bloco G-90, *campus* da Universidade Estadual de Maringá.

AGRADECIMENTOS

Agradeço a toda a equipe do Nupélia, pelos anos de árduo trabalho de campo coletando os dados aqui utilizados. Agradeço também aos meus colegas alunos do Laboratório de Ictiologia, cujos trabalhos anteriores me ajudaram a montar a tabela de características das espécies, ao Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) pela bolsa.

Espécies ameaçadas, raras e raras funcionais de peixes da planície de inundação do alto rio Paraná: perspectivas ecológicas e conservação

RESUMO

Entender como a perda de espécies está relacionada com o papel funcional em uma comunidade ou ecossistema é essencial. Explicar o porquê algumas espécies são mais suscetíveis a estarem ameaçadas do que outras é um objetivo da biologia da conservação. Além da perda de espécies, a perda de processos ecológicos que sustentam o ecossistema funcionando, pode ser a maior seqüela. Para avaliar as espécies taxonomicamente raras, raras funcionais e ameaçadas, foram usados dados dos peixes coletados entre 2000 e 2019, através de redes de arrasto com tamanho de malhas variáveis, em seis pontos amostrais da planície de inundação do alto rio Paraná. Foi considerada a abundância das espécies por ponto de coleta, sendo que para o índice de raridade taxonômica foi necessário obter o alcance geográfico, especificidade de *habitat* e tamanho da população. Para estimar a raridade funcional foram usados onze traços funcionais para a composição de uma matrix de distância, junto com uma matriz de abundância relativa para obter “*Functional Uniqueness*” e “*Functional Restrictedness*”. O resultado foi uma lista de 58 espécies. 76% destas eram simultaneamente raras taxonomicamente e funcionalmente, quatro eram ameaçadas e 17 espécies eram exóticas para a região.

Palavras-chave: raridade taxonômica; raridade funcional; conservação de espécies; peixes de água doce.

Threatened, rare, and functionally rare species of fish in the Upper Paraná River floodplain: ecological perspectives and conservation

ABSTRACT

Understanding how the loss of a species is related to its functional role in a community or ecosystem is essential. To explain why some species are more prone to be endangered than others is a major goal of conservation biology. Beyond the loss of a species, the loss of ecological processes that sustain ecosystem functioning can be the most critical influence. To evaluate the threatened, taxonomical, and functional rare species, we used fish data from 2000 to 2019, collected from nine sampling points on the Upper Paraná River floodplain (the large rivers on the floodplain, Ivinheima, Baía and Paraná, and six lakes eventually connected to the rivers: Ventura and Patos lake (Ivinheima River), Guaraná and Fechada Lakes (Baía River), Ressaco do Pau Velho and Garças Lake (Paraná River). Fish were collected using gillnets with different size mesh, and the number of individuals caught as an estimation of abundance. We obtained geographic range, habitat specificity, and population size for the taxonomical rarity index. To estimate the functional rarity, we used eleven functional traits in the composition of a functional distance matrix, together with and relative abundance matrix to obtain Functional Uniqueness and Functional Restrictedness. The outcome was a 58 species list. 76% of the species shared taxonomical and functional rarity, four were endangered, and 17 species are considered exotic to the Upper Paraná River.

Keywords: taxonomical rarity; functional rarity; species conservation; freshwater fish.

SUMMARY

1	INTRODUCTION	9
2	METHODS	11
2.1	Study area	11
2.2	Sampling	12
2.3	Species sorting and analysis	13
3	RESULTS	15
4	DISCUSSION	18
	REFERENCES	21
	APPENDIX A - Taxa filtered by sampling methodology in which we applied the rarity index. the result, rr. in crescent order. Spp = species, gri = geographic range index, hsi= habitat specificity index, psi= population size index, rr= rarity index.	27
	APPENDIX B - Fish species whose functional traits were available.	31
	APPENDIX C - Functional rarity index results.....	38

1 INTRODUCTION

Species dispersal and environmental heterogeneity play fundamental roles in the spatiotemporal biodiversity patterns (Herberich et al., 2023; Xia et al., 2023). While species vary in their dispersal ability, environmental patches vary in their level of isolation or connectedness and quality. Under low dispersal rates, local species interactions and other niche processes prevail, increasing the differences in composition between local communities when patches are heterogeneous (Mouquet & Loreau, 2002, 2003; Penha et al., 2017).

The primary selection of fish life strategies is the river dynamics, related to hydrological cycles of drought and flood, and continuous water flow (Agostinho et al., 2004; Röpke et al., 2017). The physicochemical alterations promoted by floods cause the biota to respond by morphological, anatomical, physiological, phenological, and/or ethological adaptations and produce characteristic community structures. Therefore, flooding causes perceptible alterations in biota and that biota display a defined reaction to flooding (Junk, 1989), favoring dispersal.

Rarity (here, taxonomic rarity) is widely used to assess extinction risk for conservation purposes. Rabinowitz (1981) proposed that rare species in a community could emerge from the combination of three parameters: geographic range, habitat specificity, and population size. The combination of these three characteristics defines seven forms of species rarity, with the rarest species having a small range, a high level of habitat specificity, and locally low abundance.

Concerning rare species, there is a common belief in the ‘mass ratio hypotheses’ (Grime, 1998). Traditionally using total biomass or productivity as a proxy for ecosystem functioning, where dominant species have strong effects while rare species have marginal influence (Shugart, 1984., Pastor & Post, 1986., Huston & Smith, 1986., Sala et al. 1996., Grime, 1997) However, the need to deal with ecosystem multifunctionality, resilience and/or resistance across time, and disturbances or dependence upon some keystone species, challenges this simplistic view (Power et al., 1996; Lyons et al., 2005; Violle et al., 2017).

Beyond the loss of species, there is a growing awareness that the loss of ecological processes that sustain ecosystem functioning can be the most critical outcome under accelerating global changes (Naeem et al., 2012). Rare species perform different functions in ecosystems, some being redundant with those of many other rare and common species, while others are unique (Violle et al., 2017). More recently, Violle et al. (2017), extended the scope

of Rabinowitz's (1981) classification of rarity to incorporate further differences in functional traits among taxa, defining a new component – functional rarity. Better characterizing functional rarity goes beyond the issue of understanding why species are rare or common; it can also be key to better understanding the relationship between biodiversity and ecosystem functioning. Growing consensus suggests that biodiversity-ecosystem functioning (BEF) relationships are driven by the diversity of functions carried out by species and their individuals within an ecosystem (Hooper et al., 2005; Laughlin, 2014; Cadotte et al., 2015).

One major conservation tool at the species level is the Red List of the International Union for Conservation of Nature (IUCN), whose criteria to label the extinction risks, as vulnerable, endangered, and critically endangered, take into consideration: A) the decline rate of a species, estimating the current population size comparing with the past and projecting into the future; B) small range area and decline, when its geographical range is very restricted and when other factors suggest that it is at risk; C) small population size and decline, focuses on population that are numerically small and in continuing decline; D) very small population size, allows species to be listed as threatened without evidence that there has been, is, or will be a decline of some sort; E) unfavorable quantitative analysis, allows to use any kind of quantitative analysis for assessing the risk of extinction, which is compared with the extinction-risk thresholds within given time frame (Mace et al., 2008). Since the beginning of the century, the thematic of the loss of rare species in relation to the ecosystem has been focused on, and more recently the addition of functional traits to assess ecological data (Lyons et al., 2005; Hooper et al., 2005; Naeem et al., 2013; Cadotte et al., 2015; Leitão et al., 2016; Maciel, 2021;). Works exploring the functional rarity of fish are few, and are quite recent (Toussaint et al., 2016; Grenié et al., 2018; McLean et al., 2019; Fried et al., 2021; Murgier et al., 2021; Tóth et al., 2022; Trindade-Santos et al., 2022).

Given these aspects, this study aims to evaluate the taxonomically rare, the functionally rare, as well the threatened species. Considering the key species filtered in those criteria, thus, to comprehend the response of the more fragile species in front of an environment highly regulated by dams is essential for conservation purposes, and using this work as a tool to make decisions on when, where, and how to respond to threats affecting species, ecosystems, and the services they provide in a more centered manner.

2 METHODS

2.1 Study area

The Paraná River is the second longest river in South America (4.695 km), the 10th largest river in the world for water discharge, and the fourth in drainage area (2.8×10^6 km²) (Agostinho et al., 2008). The Upper Paraná River comprises approximately the first third of the Paraná River basin, and it lies completely within the Brazilian territory. The Upper Paraná River has a drainage area of 891,000 km², representing 10.5% of the total area (Agostinho et al., 2008).

The Upper Paraná River floodplain is a stretch of 480 km between the cities of Três Lagoas, Mato Grosso do Sul State, to Guaíra, Paraná State. After the closure of the Porto Primavera Dam in 1998, the floodplain (western margin of the Paraná River) was reduced to 230 km from the Porto Primavera Dam to the Upper part of the Itaipu Reservoir, which is one of the last stretches of the Paraná River with running waters. This stretch has high habitat heterogeneity and still maintains a great biodiversity of terrestrial and aquatic organisms (Agostinho et al., 2004a.; Agostinho et al., 2004b).

Despite the ecological importance of the floodplain, the Paraná River and its tributaries have more than 145 large dams, most of them in cascade. These reservoirs caused alterations in the annual flood regime of the Paraná River and its floodplain (Agostinho et al., 2005.; Agostinho et al., 2004; Sanches et al., 2006) but the flood pulse is still the main force function for the biota of the region. The flood period in the Paraná River usually occurs from November/December to April/May, while the low-water period coincides with the winter, between June and September. The occurrence of two or three annual flood pulses is quite common during periods of high water levels (Agostinho et al., 2009; Lima et al., 2017).

In the floodplain, the river has a wide-braided channel with a low slope and heavy accumulation of sediments on its bed, creating sandbanks and islands. There are also secondary but equally important channels, the Baía and Ivinheima rivers, and the lower courses of rivers entering on the right bank (e.g. Amambai, and Iguatemi rivers) forming a complex anastomosis in this stretch. The left margin of the river is steeper, and its tributaries have larger slopes (Paranapema, Ivai, and Piquiri rivers) with smaller floodplains (Figure 1).




Figure 1. Map of the study area. 1= Ventura Lagoon, 2= Patos Lagoon, 3= Ivinhema River, 4= Guaraná Lagoon, 5= Baía River, 6= Fechada Lagoon, 7= Pau Vêio Backwater, 8= Paraná River, 9= Garças Lagoon. Fonte: Andreotti et al. (2021)

2.2 Sampling

Sampling was conducted quarterly, from March 2000 until December 2019, in nine sampling sites, as follow: three including the large rivers on the floodplain, Ivinhema, Baía and Paraná, and six sampling sites on floodplain lakes connected or not to the rivers: Ventura and Patos Lagoon (Ivinhema River), Guaraná and Fechada Lagoon (Baía River), Ressaco do Pau Velho and Garças Lagoon (Paraná River) (Figure 1).

Fishes were collected using 20 m-long and 1.70 m-high gillnets with different mesh sizes (2.4 to 16 cm between opposite knots) that were exposed for 24 h and checked every 8 h, 16 h, and 22 h. Captured fish were anesthetized with 5% benzocaine and euthanized. All captured fish were identified according to Garavello (1979), Garavello and Britski (1987, 1988, 1990), Britski *et al.* (2007), Graça and Pavanelli (2002), and Ota *et al.* (2018). Because the fishing effort was the same for all samplings, I considered the number of individuals caught as an estimation of abundance.

2.3 Species sorting and analysis

For the definition of endangered species, we considered the Red Book of Threatened Brazilian Fauna: Vol. VI – Fishes (ICMBio, 2018), Portaria MMA 148/ 2022 and the IUCN Red List of Threatened Species (IUCN, 2022).

The taxonomic rare species were selected using a package in the R environment (R CoreTeam, 2023) introduced by Maciel (2021) called “*rrindex*”. The idea behind the index follows the reasoning proposed by Rabinowitz (1981). The output of this package is a matrix with five columns, containing the following parameters for each species: species (*spp*), the geographic range index (*gri*), habitat specificity index (*hsi*), the population size index (*psi*), and an average among them, the taxonomic rarity index (*rr*). The “*rrindex* package” requires a series of inputs. The first one is a list with the name of the species as a factor, and the name of each species can appear in the list more than once. The second and third parameters are numeric vectors containing the information for latitude and longitude where species were registered. The fourth input is of the string type and represents the abundance of the species in a given area. The fifth input is a factor containing information on the type of habitat in which the species was collected. In this case, we used “*rivi*”, “*rpar*”, “*rbai*”, to address each river and its two respective lakes temporally connected.

To quantify functional rarity, an R package named “*funrar*” was used, based on occurrence and trait data (R Core Team, 2023; Grenié et al., 2017). This package computes Functional Distinctiveness (*Di*), Functional Uniqueness (*Ui*), Taxon Scarcity (*Si*), and Taxon Restrictedness (*Ri*) using a species-by-site or community composition matrix with either the presence-absence or relative abundance of species, and a functional dissimilarity matrix. As emphasized by Violle et al. (2017), a species can be functionally distinct in a given community but not functionally unique in an entire region. In this regard, distinctiveness and scarcity are used to uncover local rarity; uniqueness and restrictedness are used regionally. In this work, I considered the floodplain on a regional scale. That is, we considered only uniqueness (*Ui*), and restrictedness (*Ri*). In addition, after applying these functions, I used a Spearman’s correlation to check the relationship between these two indices separately. A Principal Coordinates Analysis (PCoA) was carried out to better visualize the functional dissimilarity of the species.

At the regional scale, Functional Uniqueness is relevant to assess the role of functional originality. It can represent how taxa depart from a regional pool due to specific biogeographical and evolutionary legacies and the index result varies from 0 to 1. Taxon restrictedness is an index used between 0 and 1. It tends to one when a species is present in fewer sites of the species-by-site matrix. To obtain the functional rarity, the average between U_i e R_i is calculated.

The functional traits used in this work were previously obtained through the works of Rodrigues et al. (2020) and Baumgartner et al. (2018), and they were considered to better characterize the species' functional role and response regarding the floodplain environment (Appendix B) (Winemiller et al., 2015; Villéger et al., 2017).

They are as follows:

Trophic category	detritivore, insectivore, piscivore, herbivore, omnivore, and planktivore (according to Hahn et al.,2004)
Position in the water column	benthopelagic, pelagic, and demersal (according to Graça e Pavanelli, 2007)
Body shape	fusiform, deep, laterally compressed, cylindrical, flattened, anguilliform (according to Graça e Pavanelli, 2007)
Crypsis	low, moderate, and extreme (according to Graça & Pavanelli, 2007; Calegari et al.,2013; Souza-Filho & Shibatta,2007. Deprá et al.,2015; Loboda et al.2015)
Standard length	(according to Costa & Soares,2015; Graça & Pavanelli, 2007; Deprá et al.,2015; Loboda et al.2015)
Eye position	lateral, dorsolateral, superior and inferior (according to Britski et al.,2007; Roberts, 2015; Varella,2011)
Mouth position	subterminal, terminal, inferior, superior, and lateral (according to Graça & Pavanelli, 2007; Castro et al.,20014; Britski et al.,2007; Marinho et al.,2008; Zarske et al.2008)
Diel activity	crepuscular, nocturnal and diurnal (according to Britski et al.,2007; Reis et al., 2003; Nakatani et al.,2001)
Fecundation and parental care	external fecundation with parental care, external fecundation without parental care, and internal fecundation (according to Nakatani et al., 2003; Suzuki et al., 2005; Graça & Pavanelli, 2007; Froese & Pauly, 2015)
Spawning	parceled, total (according to Nakatani et al., 2001; Suzuki et al., 2005; Graça & Pavanelli, 2007; Froese & Pauly, 2015)
Migration	migratory, sedentary (according to Graça & Pavanelli, 2007; Froese & Pauly, 2015)

For a more accurate result and concerning conservational purposes, only the species correctly identified were considered on the final list.

3 RESULTS

Four species matched one or more criteria established to define threatened species: *Aphyocheirodon hemigrammus*, *Brycon orbignyianus*, *Myloplus tiete*, and *Pseudoplatystoma corruscans*. Using the rarity metrics (Figure 2), the number of rare species found was 51 (33,5% of total species) (Appendix A). For they presented *rr* above average ($rr = 0.36$) measured for all species ($n = 152$).




Figure 2. Boxplot showing the result of the rarity metrics for geographic range, habitat specificity, population size indices, and the average between them. (gri) = geographic range index, (hsi) = habitat specificity index, (psi) = population size index, and (rr) = rarity index

Among the 152 fish species registered, 127 species had trait information available. For each species, I calculated Functional Uniqueness and Taxon Restrictedness and averaged them by each species. Out of the 127 species, 44% (56) species were above the average (0.28) of functional rarity (Appendix C), but even so, the index for the rarest functionally species was only 0.66, *Farlowella hahni*. Despite of this value, the Functional Restrictedness for this species was 0.88, indicating that it occurred in a few sampling sites, and the Functional Uniqueness was 0.44, which means that its functional traits are shared with some other few species. In general, most species are functionally redundant, with low values of Functional Uniqueness, but 33% (42) of them are geographically restricted, present in just one sampling site (Figure 3). At the species level, Functional Uniqueness and Taxon Restrictedness were weakly inversely correlated (Spearman's $\rho = -0.10$, $p = 0.23$). To better visualize the functional distance in contrast to the Functional Rarity Index, I made a PCOA. In which the position of the species is located differently than the index (Figure 4).




Figure 3. Taxon Restrictedness against Functional Uniqueness per species.




Figure 4. PCOA species scores showing the functional distance of the species.

Matching at least one of the three criteria, 58 species were pointed out (Table 1). For the species *Aphyocheirodon hemigrammus*, no functional trait was available. So, it is the only one without the metrics for functional and taxonomical rarity. 76% of the listed species (n=58, excluding the one cited above) shared taxonomical and functional rarity. 17 species are considered exotic to the Upper Paraná River according to Ota et al. (2018).

Table 1. Fish species and the respective criteria in which they fit. EN= endangered, VU= vulnerable, CR= critically endangered. * Alien species to the Upper Paraná River.

Species	Taxonomic Rarity Index	Functional Rarity Index	IUCN	Red Book	Portaria MMA Nº 148,07.07.22
<i>Acestrorhynchus pantaneiro</i> Menezes, 1992*	0,36	31			
<i>Aequidens plagiozonatus</i> Kullander, 1984	0,73	0,45			
<i>Apareiodon affinis</i> (Steindachner, 1879)	0,49	0,51			
<i>Aphyocheirodon hemigrammus</i> Eigenmann, 1915			VU	VU	
<i>Apteronotus caudimaculosus</i> Santana, 2003	1	0,44			
<i>Apteronotus ellisi</i> (Arámburu, 1957)	0,69	0,44			
<i>Brachyhyopomus gauderio</i> Giora, Malabarba, 2009*	1	0,44			
<i>Brycon hilarii</i> (Valenciennes, 1850)*	0,43	0,47			
<i>Brycon orbignyanus</i> (Valenciennes, 1850)	0,16	0,13		EN	CR
<i>Catathyridium jenynsii</i> (Günther, 1862)*	0,19	0,31			
<i>Colossoma macropomum</i> (Cuvier, 1818)*	0,83	0,47			
<i>Crenicichla haroldoi</i> Luengo, Britski, 1974	0,83	0,5			
<i>Crenicichla jaguarensis</i> Haseman, 1911	0,44	0,39			
<i>Crenicichla nierderleinii</i> Varella (2011)	0,77	0,59			
<i>Eigenmannia virescens</i> (Valenciennes, 1836)	0,51	0,4			
<i>Erythrinus erythrinus</i> (Bloch, Schneider, 1801)*	0,21	0,4			
<i>Farlowella hahni</i> Meinken, 1937*	1	0,66			
<i>Galeocharax gulo</i> (Cope, 1870)	0,21	0,42			
<i>Gymnotus pantanal</i> Fernandes, Albert, Daniel-Silva, Lopes, Crampton, Almeida-Toledo, 2005	0,22	0,31			
<i>Gymnotus paraguensis</i> Albert, Crampton, 2003	0,27	0,37			
<i>Hoplias intermedius</i> (Günther, 1864)	1	0,48			
<i>Hypostomus albopunctatus</i> (Regan, 1908)	0,55	0,43			
<i>Hypostomus cf. strigaticeps</i> (Regan, 1908)	0,23	0,37			
<i>Hypostomus commersoni</i> Valenciennes, 1836*	0,54	0,43			
<i>Hypostomus hermanni</i> (Ihering, 1905)	0,76	0,43			
<i>Hypostomus microstomus</i> Weber, 1987*	1	0,48			
<i>Hypostomus ternetzi</i> (Boulenger, 1895)*	1	0,48			
<i>Laetacara araguaiae</i> Ottoni, Costa, 2009*	0,65	0,38			
<i>Leporellus vittatus</i> (Valenciennes, 1850)	0,7	0,53			
<i>Leporinus octofasciatus</i> Steindachner, 1915	0,66	0,47			
<i>Leporinus striatus</i> Kner, 1858	0,54	0,48			
<i>Loricaria prolixa</i> Isbrücker, Nijssen, 1978	0,24	0,42			
<i>Megalancistrus parananus</i> (Peters, 1881)	0,27	0,42			
<i>Megalonema platanum</i> (Günther, 1880)	0,66	0,51			
<i>Moenkhausia bonita</i> Benine, Castro, Sabino, 2004	1	0,53			
<i>Moenkhausia cf. gracilima</i> Eigenmann, 1908	1	0,53			
<i>Moenkhausia forestii</i> Benine, Mariguela, Oliveira, 2009*	1	0,53			
<i>Myloplus tiete</i> (Eigenmann, Norris, 1900)	0,57	0,43			EN
<i>Oligosarcus pinto</i> Campos, 1945	0,75	0,58			
<i>Ossancora eigenmanni</i> (Boulenger, 1895)*	0,15	0,29			
<i>Parodon nasus</i> Kner, 1859	0,52	0,57			
<i>Piabarchus stramineus</i> (Eigenmann, 1908)	0,29	0,46			
<i>Pimelodella taenioptera</i> Miranda-Ribeiro, 1914*	0,54	0,47			
<i>Pimelodus microstoma</i> Steindachner, 1877	0,2	0,32			
<i>Pinirampus pirinampu</i> (Agassiz, 1829)	0,22	0,41			
<i>Platydoras armatulus</i> (Valenciennes, 1840)*	0,24	0,37			
<i>Psalidodon paranae</i> (Eigenmann, 1914)	0,83	0,44			
<i>Psalidodon schubarti</i> *	0,51	0,33			
<i>Psellogrammus kennedyi</i> (Eigenmann, 1903)*	0,12	0,28			

<i>Pseudopimelodus mangurus</i> (Valenciennes, 1835)	0,56	0,57			
<i>Pseudoplatystoma corruscans</i> (Spix, Agassiz, 1829)	0,12	0,09			VU
<i>Pseudoplatystoma reticulatum</i> Eigenmann, 1889*	0,5	0,47			
<i>Rhamdia quelen</i> (Quoy, Gaimard, 1824)	0,23	0,4			
<i>Rhinodoras dorbignyi</i> Kner, 1855)	0,32	0,48			
<i>Schizodon nasutus</i> Kner, 1858	0,12	0,4			
<i>Synbranchus marmoratus</i> Bloch, 1795	0,83	0,61			
<i>Triportheus nematurus</i> (Kner, 1858)*	0,68	0,54			
<i>Zungaro jahu</i> (Ihering, 1898)	0,51	0,52			

4 DISCUSSION

Brycon orbignyanus was the most threatened species on the list (MMA,2022). And even so, it did not appear in any of the rarity indexes. Fact that can be explained since I used only the floodplain as a source of data. It is a migratory species, and the main threat is pollution, followed by the destruction of riparian vegetation, silting of rivers and dams in cascade. (ICMBio, 2018; Agostinho and Gomes, 2004). This species is very sensitive to environmental changes and requires an area with preserved riparian forests for food and consolidation of the reproductive process (Cecílio and Agostinho, 1997). With a loss of 97% of its extent of occurrence area, nowadays in Brazil, the Upper Paraná River (including its floodplain and some tributaries) is the last natural habitat for the species (ICMBio, 2018). Another migratory threatened species on the list is *Pseudoplatystoma corruscans* (VU-MMA, 2022). Whose main threats are damming, overfishing, and silting (Mello et al., 2009; Silva, 2015). *Aphyocheirodon hemigrammus* is endemic to the Upper Paraná River Basin. Found in low abundance in lentic water bodies, preferring marginal lakes (ICMBio,2018). *Myloplus tiete* is a riverine species, and its threat is related to the expansion of hydropower dams and introduction of invasive species such as *Cichla* spp (peacock bass) (Magalhães et al., 2019).

Rare species have become the focal species for conservation in the field of biodiversity conservation. In this sense, a simple approach like the “rrindex” used in this work can reduce the gap between science and decision-making. Another challenge is to try to understand how rarity can be associated with the risk of extinction. These questions have already been addressed using geographic range or population size as response variables. The index used here provides the ability to test this and other questions from the synthesis of three parameters (geographic

range, habitat specificity, and population size) that are already recognized as important aspects of classifying rare species in a community of different taxa (Maciel, 2021).

At regional scale, Functional Uniqueness can represent how taxa depart from a regional pool due to specific biogeographical and evolutionary legacies and should then be estimated based on the whole site-species matrix (Grenié et al., 2017).

Some species such as *Farlowella hahni*, are non-native, which occurrence can be associated with the filling of the Itaipu Reservoir and the consequent inundation of the Sete Quedas Falls (Ota et al. 2018). Therefore, despite its uniqueness, in the end, it doesn't match the conservational idea behind this work.

In general, except for *Farlowella hahni*, no other species in the final list has stood out from another. Whether by a high value for taxonomical rarity (max=0.83, *Synbranchus marmoratus*, *Psalidodon paranae*, and *Crenicichla haroldoi*), or high value for functional rarity (max= 0.61, *Synbranchus marmoratus*). It might be explained by the sampling methodology, being captured accidentally by a device not focused on fish of their size, morphology, life habits, and the redundancy of functions they possess amongst themselves. Knowing specifically the causes of rarity can predict in greater success the efforts for species preservation. Determining a boundary where rarity begins is a theme that has been already discussed by many authors (Dajoz, 1983., Maitland, 1985., Olden et al. 2008., Leitão et al. 2016). As well as the fluctuation, similarity, diversity, and turnover (Cadotte et al., 2015; Baumgartner et al., 2018; Cantarute et al., 2018) of functional traits in some aspects, quite similar and the building blocks of the indexes I worked with here, I hoped that it would become clear the soft transition of one species to another considering the functional traits and the multi-factor aspect that encompasses the traditional taxonomic rarity. If rare species mainly support roles that are also played by common species, then we may expect mild consequences following their extinction. On the other hand, if rare species over-contribute to functional structure, then their extinction may lead to a harsh loss of ecological processes, as demonstrated by Pendleton et al. (2014) and Leitão et al. (2016) which showed that the loss of rare species may reduce the functional richness, specialization, and originality of assemblages more than expected under a random loss of species.

Progress in understanding how the loss of a species is related to its functional role in a community or ecosystem is essential. Studies showing the effects of species richness on

ecosystem functioning are controversial. Ecosystem functions do not saturate at low species richness, many studies show positive effects of biodiversity on function (Oliver et al., 2015; Gonzalez et al., 2020; Moi et al., 2021), while others do not (Schwartz et al., 2000; Winfree et al., 2015). It is not yet clear how results from small-scale BEF experiments can be scaled up to larger spatial scales relevant to conservation. Although, realistic patterns of species extinction can result in BEF effects different from those predicted by random-loss experiments (Schwartz et al., 2000; Srivastava & Vellend, 2005).

REFERENCES

- Abell, R., M.L., Thieme, C. Revenga, M. Bryer, M. Kottelat & N. Bogutskaya, 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *Bioscience* 58, 403–414.
- Agostinho A.A., S.M. Thomaz, C.V. Minte-Vera & K.O. Winemiller, 2000. Biodiversity in the high Paraná River floodplain. In B. Gopal, W.J. Junk, J.A. Davis eds. *Wetlands Biodiversity*. pp. 89-118. Leiden, The Netherlands, Backhuys Publishers. 2000.
- Agostinho A.A., L.C. Gomes & F. Pelicice, 2008. Dams and the fish fauna of the Neotropical region: Impacts and management related to diversity and fisheries. *Braz. J. Biol.*, 68(4, Suppl.): 1119-1132, 2008.
- Agostinho A.A., L.C. Gomes & M. Zalewski, 2001. The importance of floodplains for the dynamics of fish communities of the upper river Paraná. *Ecohydrology & Hydrobiology*, 1: 209-217.
- Agostinho, A. A., F.M. Pelicice, & L.C. Gomes, 2008. Dams and the fish fauna of the Neotropical region: impacts and management related to diversity and fisheries. *Brazilian Journal of Biology* 68: 1119–1132.
- Agostinho, A. A., L. C. Gomes, S. Veríssimo, & E.K. Okada, 2004. Flood regime, dam regulation and fish in the Upper Paraná River: effects on assemblage attributes, reproduction and recruitment. *Rev. Fish Biol. Fish.* 14, 11–19.
- Agostinho, A.A., C. Bonecker, & L.C. Gomes, 2009. Effects of water quantity on connectivity: The case of the upper Paraná River floodplain. *Ecohydrology and Hydrobiology Elsevier B.V.* 9: 99–113.
- Agostinho, A.A., S.M. Thomaz, & L.C. Gomes, 2004-Threats for biodiversity regulated by dams. *Ecohydrology and Hydrobiology*. V.4 N.3: 255-256
- Andreotti, G.F., J.C. Alves, D.C. Alves, A.A. Angelo & L.C. Gomes, 2021. The response of fish functional diversity to EL Niño Southern Oscillation (ENSO) in a Neotropical Floodplain. *Hydrobiologia*. 848:1207–1218.
- Baumgartner, M. T., A. G. de Oliveira, A. A. Agostinho, & L. C. Gomes, 2018. Fish functional diversity responses following flood pulses in the upper Paraná River floodplain. *Ecology of Freshwater Fish Blackwell Munksgaard* 27: 910–919.
- Britski H.A., K.Z.S. Silimon, 2007. Lopes BS. *Peixes do Pantanal: manual de identificação*. Brasília, DF: Embrapa Pantanal.
- Cadotte, M. W., C. A. Arnillas, S. W. Livingstone, & S. L. E. Yasui, 2015a. Predicting communities from functional traits. *Trends in Ecology and Evolution*. Elsevier Ltd, 510–511.
- Calegari, B. B., P. Lehmann & R.E. Reis, 2013. Two new species of cascudinhos of the genus *Otothyropsis* (Siluriformes: Hypoptopomatinae) from the rio Paraná basin, Brazil. *Zootaxa*, 3619: 130-144.
- Cecílio, E.B, A.A. Agostinho, H.F. Júlio Jr. & C.S. Pavanelli, 1997. Colonização ictiofaunística do reservatório de Itaipu e sua área adjacentes. *Revista Brasileira de Zoologia*, CURITIBA-PR, v. 14, p. 1-14.

- Comunello, E., E.E. Souza-Filho, P. C. Rocha & M. Nanni, 2003. "Dinâmica de inundação de áreas sazonalmente alagáveis na planície aluvial do Alto Rio Paraná: estudo preliminar,". in Anais do XI Congresso Brasileiro de Sensoriamento Remoto (Belo Horizonte: INPE), 2459–2466.
- Dajoz, R, 1983. Ecologia Geral, 3.ed. Petrópolis: Vozes. 474p.
- Deprá, G.C., R.R.Ota, F. Souza, W.J. Graça & C.S. Pavanelli, 2015. Widening the geographical distribution of *Pimelodus mysteriosus* Azpelicueta 1998 (Siluriformes: Pimelodidae) to the upper Paraná River, with diagnosis for syntopic congeners. Biota Neotropica. 15: 1–6.
- Froese, R. & D. Pauly, 2021. Fishbase. www.fishbase.org. Accessed November 2023
- Garavello J.C. & H.A., Britski, 1987. Duas novas espécies do gênero *Leporinus* Spix, 1829, da bacia o alto rio Paraná (Teleostei, Anostomidae). Com Mus Cien PUCRS.44:153-65.
- Garavello J.C. & H.A., Britski, 1988. *Leporinus macrocephalus* sp. n. da bacia do rio Paraguai (Ostariophysi, Anostomidae). Naturalia.13:67-74.
- Garavello J.C. & H.A., Britski, 1990. Duas novas espécies do gênero *Schizodon* Agassiz da bacia do alto Paraná, Brasil, América do Sul (Ostariophysi, Anostomidae). Naturalia. 15:153-70.
- Garavello J.C.,1979. Revisão taxonômica do gênero *Leporinus* Spix, 1829 (Ostariophysi, Anostomidae). [PhD Thesis]. São Paulo, SP: Universidade de São Paulo.
- Gonzalez, A., R. M. Germain, D. S. Srivastava, E. Filotas, L. E. Dee, D. Gravel, P. L. Thompson, F. Isbell, S. Wang, S. Kéfi, J. Montoya, Y. R. Zelnik, & M. Loreau, 2020. Scaling-up biodiversity-ecosystem functioning research. Ecology Letters Blackwell Publishing Ltd 23: 757–776.
- Graça, W.J., C.S. Pavanelli, 2007. Peixes da planície de inundação do alto rio Paraná e áreas adjacentes. 1. ed. Maringá: EDUEM,1. 241
- Grenié, M., D. Mouillot, S. Villéger, P. Denelle, C. M. Tucker, F. Munoz, & C. Violle, 2018. Functional rarity of coral reef fishes at the global scale: Hotspots and challenges for conservation. Biological Conservation Elsevier Ltd 226: 288–299.
- Grenié, M., P. Danelle, C.M. Tucker, F. Munoz & C. Violle, 2018. Functional rarity of coral reef fishes at the global scale: hotspots and challenges for conservation. Biol. Conserv. 226, 288–299.
- Grenié, M., P. Denelle, C. M. Tucker, F. Munoz, & C. Violle, 2017. funrar: An R package to characterize functional rarity. Diversity and Distributions. Blackwell Publishing Ltd, 1365–1371.
- Grime, J.P., 1998. Benefits of plant diversity to ecosystems: immediate, filter and founder effects. J. Ecol., 86, 902–910.
- Hahn, N. S., R. Fugi, & I.F. Andrian, 2004. Trophic ecology of the fish assemblages. In: The Upper Paraná River and its Floodplain: Physical aspects, Ecology and Conservation, pp. 247-269.
- Herberich, M.H., Sebastian G., Katja T., 2023. Environmental heterogeneity promotes coexistence among plant life-history strategies through stabilizing mechanisms in space and time, Basic and Applied Ecology, 71:45-56. <https://doi.org/10.1016/j.baae.2023.05.001>
- Hooper, D. U., F. S. Chapin, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naeem, B. Schmid, H. Seta"la", S. Seta"la, S. Seta"la", A. J. Symstad, J. Vandermeer, & D. A. Wardle, 2005. ESA Report Effects of Biodiversity on Ecosystem Functioning: A Consensus of Current Knowledge. Ecological Monographs.75,3-35.

- Huston, M.A & Smith, T.M., 1987. Plant succession: life history and competition. *American Naturalist*, 130,168-198.
- Instituto Chico Mendes de Conservação da Biodiversidade – ICMBio, 2018. Livro Vermelho da Fauna Brasileira Ameaçada de Extinção: Volume VI – Peixes. Brasília, DF.7.
- IUCN. 2023. The IUCN Red List of Threatened Species. Version 2023-1. <https://www.iucnredlist.org>. Accessed on [20.05.2023].
- Jeppesen, E., T. Mehner, I.J. Winfield., et al., 2012. Impacts of climate warming on the long-term dynamics of key fish species in 24 European lakes. *Hydrobiologia* 694:1–39 DOI 10.1007/s10750-012-1182-1
- Junk, W.J, 1989. The Flood Pulse concept in River-Floodplain System. *Canadian Journal of Fisheries and Aquatic Sciences*, Canada. 106: 110-127,
- Laughlin, D. C., 2014. Applying trait-based models to achieve functional targets for theory-driven ecological restoration. *Ecology Letters Blackwell Publishing Ltd* 17: 771–784.
- Laughlin, D.C,2014. Applying trait-based models to achieve functional targets for theory-driven ecological restoration. *Ecol. Lett.* 17, 771–784 26.
- Leitão, R. P., J. Zuanon, S. Villéger, S. E. Williams, C. Baraloto, C. Fortune, F. P. Mendonça, & D. Mouillot, 2016. Rare species contribute disproportionately to the functional structure of species assemblages. *Proceedings of the Royal Society B: Biological Sciences Royal Society of London*.283.
- Lima, C., A. Aghakouchak, & U. Lall, 2017. Classification of mechanisms, Climatic Context, Areal Scaling, and Synchronization of floods: the hydroclimatology of floods in the Upper Paraná River Basin, Brazil. *Earth Syst. Dynam. Discuss* .
- Loboda, T.S. & M.R. de Carvalho, 2013. Systematic revision of the *Potamotrygon motoro* (Müller & Henle, 1841) species complex in the Paraná-Paraguay basin, with description of two new ocellated species (Chondrichthyes: Myliobatiformes: Potamotrygonidae). *Neotrop. Ichthyol/* 11(4):693-737.
- Lyons, K. G., C. A. Brigham, B. H. Traut, & M. W. Schwartz, 2005. Rare species and ecosystem functioning. *Conservation Biology* . , 1019–1024.
- Mace, G. M., N. J. Collar, K. J. Gaston, C. Hilton-Taylor, H. R. Akçakaya, N. Leader-Williams, E. J. Milner-Gulland, & S. N. Stuart, 2008. Quantification of extinction risk: IUCN’s system for classifying threatened species. *Conservation Biology*. 1424–1442.
- Maciel, E. A., 2021. An index for assessing the rare species of a community. *Ecological Indicators Elsevier B.V.* 124.
- Magalhães, A.L.B., F.M. Pelicice & D.P. Lima-Junior, 2019. Riscos ambientais e socioeconômicos do Projeto de Lei que visa a proteção de espécies invasoras (tucunaré azul e tucunaré amarelo) no Estado do Paraná. (Nota Técnica).
- Maitland, P. S,1985. Criteria for the selection of important sites for freshwater fish in the British Isles. *Biological Conservation*, 31: 335-353
- Marinho, M.M.F., F.R. Carvalho, F. Langeani & F.L. Tatsumi, 2008. A new *Hemigrammus* Gill from upper rio Paraná system, Southeastern Brazil (Characiformes: Characidae). *Zootaxa* 1724:52-60

- Moi, D., F. Lansac-Tôha, G. Romero, P. Kratina, D. Perkins, F. Teixeira De Mello, & J. Heino, 2021. Human pressure drives biodiversity-multifunctionality relationships in neotropical wetlands. *Nature Ecology & Evolution*, 1-18. <https://doi.org/10.1038/s41559-022-01827-7>
- Moi, D.A., J. Ernandes-Silva, M.T. Baumgartner & R.P. Mormul, 2019. The effects of river-level oscillations on the macroinvertebrate community in a river–floodplain system. *Limnology* 21:219–232.
- Mormul, R. P., S. M. Thomaz, A.A. Agostinho, C.C. Bonecker, & N. Mazzeo, 2012. Migratory benthic fishes may induce regime shifts in a tropical floodplain pond. *Freshw. Biol.* 57, 1592–1602. doi: 10.1111/j.1365-2427.2012.02820.x
- Mouquet, N., & M. Loreau, 2002. Coexistence in metacommunities: The regional similarity hypothesis. *American Naturalist*.420–426.
- Mouquet, N., & M. Loreau, 2003. Community Patterns in Source-Sink Metacommunities. *American Naturalist* 162: 544–557.
- Naeem, S., J.E., Duffy, & E. Zavaleta, 2012. The functions of biological diversity in an age of extinction. *Science* 336, 1401–1406.
- Nakatani, K., A.A. Agostinho, G. Baumgartner, A. Bialecki, P.V. Sanches, M.C. Makrakis & C.S. Pavanelli, 2001. *Ovos e larvas de peixes de água doce: desenvolvimento e manual de identificação*.Eduem.
- Olden, J.D., N.L. Poof, K.R. Bestgen, 2008. Trait synergisms and the rarity, extirpation, and extinction risk of desert fishes. *Ecology*, 89:847–856.
- Oliver, T. H., M. S. Heard, N. J. B. Isaac, D. B. Roy, D. Procter, F. Eigenbrod, R. Freckleton, A. Hector, C. D. L. Orme, O. L. Petchey, V. Proença, D. Raffaelli, K. B. Suttle, G. M. Mace, B. Martín-López, B. A. Woodcock, & J. M. Bullock, 2015. Biodiversity and Resilience of Ecosystem Functions. *Trends in Ecology and Evolution*. Elsevier Ltd, 673–684.
- Ota, R. R., G. de C. Deprá, W. J. da Graça, & C. S. Pavanelli, 2018. *Peixes da planície de inundação do alto rio Paraná e áreas adjacentes: Revised, annotated and updated*. Neotropical Ichthyology Sociedade Brasileira de Ictiologia 16:.
- Pastor, J. & Post, W.M., 1986. Influence of climate, soil moisture and succession on forest carbon and nitrogen cycles. *Biogeochemistry*, 2, 3-27.
- Pendleton, R. M., D. J. Hoeinghaus, L. C. Gomes, & A. A. Agostinho, 2014. Loss of rare fish species from tropical floodplain food webs affects community structure and ecosystem multifunctionality in a mesocosm experiment. *PLoS ONE Public Library of Science* 9.
- Penha, J., Victor, L.L., Jean, C.G.O., Lúcia, M., 2017. Interchange between flooding and drying, and spatial connectivity control the fish metacommunity structure in lakes of the Pantanal wetland. *Hydrobiologia* 797:115-126. <https://doi.org/10.1007/s10750-017-3164-9>.
- Power, M. E., D. Tilman, J. A. Estes, B. A. Menge, W. J. Bond, L. S. Mills, G. Daily, J. C. Castilla, J. Lubchenco, & R. T. Paine, 1996. Challenges in the Quest for Keystones Identifying keystone species is difficult-but essential to understanding how loss of species will affect ecosystems. <https://academic.oup.com/bioscience/article/46/8/609/237132>.
- Rabinowitz D.,1981. Seven forms of rarity. In *Biological aspects of rare plant conservation*. Chichester, UK: John Wiley.205–217.

- Reis, R. E, 2003. Family Calliichthyidae. In: Reis, R. E., S.O. Kullander, & C.J. Ferraris Jr. Check list of freshwater fishes of South and Central America. Porto Alegre, Edipucrs, p. 291-309.
- Roberts, T. R, 2015. Mimicry of a bean seed by the Amazonian Aspredinid catfish *Amaralia hypsura* (Kner, 1855), with notes on vegetative camouflage by fishes. *Aqua International Journal of Ichthyology*, 21: 110-119.
- Rocha, P. C., M. Santos & E.E. Souza-Filho, 2001. "Alterações no regime hidrológico do alto rio Paraná como resposta ao controle de descargas efetuado por grandes barramentos a montante," in *Anais do VIII Encuentro de Geógrafos de América Latina* (Santiago), 28–39.
- Röpke, C. P., S. Amadio, J. Zuanon, E. J. G. Ferreira, C. P. De Deus, T. H. S. Pires, & K. O. Winemiller, 2017. Simultaneous abrupt shifts in hydrology and fish assemblage structure in a floodplain lake in the central Amazon. *Scientific Reports Nature Publishing Group* 7:.
- RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL <http://www.rstudio.com/>.
- Sanches, P. V., K. Nakatani, A. Bialecki, G. Baumgartner, L. C. Gomes, & E. A. Luiz, 2006. Flow regulation by Dams affecting ichthyoplankton: The case of the Porto Primavera Dam, Paraná River, Brazil. *River Research and Applications* 22: 555–565.
- Schwartz, M.W., C.A. Brigham, J.D. Hoeksema, K.G. Lyons, M.H. Mills & P.J. Van Mantgem, 2000. Linking biodiversity to ecosystem function: implications for conservation ecology. *Oecologia*, 122, 297–305.
- Shugart, H.H., 1984. *A theory of forest dynamics*. Springer Verlag, New York.
- Souza-Filho, H. S. D.; Shibatta, O. A. (2007). Descrição de uma espécie nova de *Pimelodus* (Siluriformes, Pimelodidae) da bacia do alto rio Paraguai. *Iheringia. Série Zoologia*. 97(4): 472-480
- Srivastava, D.S. & M. Vellend, 2005. Biodiversity-ecosystem function research: is it relevant to conservation? *Annual Review of Ecology, Evolution, and Systematics*. 36, 267–294.
- Tóth, R., I. Czeglédi, P. Takács, P. A. Tedesco, & T. Erős, 2022. Taxonomic rarity and functional originality of freshwater fishes and their responses to anthropogenic habitat alterations. *Aquatic Conservation: Marine and Freshwater Ecosystems John Wiley and Sons Ltd* 32: 484–494.
- Toussaint, A., N. Charpin, S. Brosse, & S. Villéger, 2016. Global functional diversity of freshwater fish is concentrated in the Neotropics while functional vulnerability is widespread. *Scientific Reports Nature Publishing Group* 6:.
- Trindade-Santos, I., F. Moyes, & A. E. Magurran, 2022. Global patterns in functional rarity of marine fish. *Nature Communications Nature Research* 13:.
- Varella, H. R, 2011. Revisão taxonômica das espécies de *Crenicichla* Heckel das bacias dos rios Paraná e Paraguai (Teleostei: Cichlidae). Dissertação de mestrado - Instituto de Biociências da Universidade de São Paulo. Departamento de Zoologia.
- Villéger, S., S. Brosse, M. Mouchet, D. Mouillot, & M. J. Vanni, 2017. Functional ecology of fish: current approaches and future challenges. *Aquatic Sciences Birkhauser Verlag AG* 79: 783–801.
- Violle, C., W. Thuiller, N. Mouquet, F. Munoz, N. J. B. Kraft, M. W. Cadotte, S. W. Livingstone, & D. Mouillot, 2017a. Functional Rarity: The Ecology of Outliers. *Trends in Ecology and Evolution. Elsevier Ltd*, 356–367.

Winemiller, K. O., D. B. Fitzgerald, L. M. Bower, & E. R. Pianka, 2015. Functional traits, convergent evolution, and periodic tables of niches. *Ecology Letters* Blackwell Publishing Ltd 18: 737–751.

Winfree, R., J. W. Fox, N. M. Williams, J. R. Reilly, & D. P. Cariveau, 2015. Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecology Letters* 18: 626–635.

Xia Z., Jani, H., Fandong, Y., Yonfeng, H., Fei, L. Jianwei, W., 2022. Spatial patterns of site and species contributions to β diversity in riverine fish assemblages, *Ecological Indicators*, 145. <https://doi.org/10.1016/j.ecolind.2022.109728>.

Zarske, A., P.-Y. Le Bail & J. Géry, 2006. New and poorly known characiform fishes from French Guiana. 1. Two new tetras of the genera *Hemigrammus* and *Hyphessobrycon* (Teleostei: Characiformes: Characidae). *Zool. Abhand. Dresden* 55:17-30.

Silva, A.B., 2015. Influência de fatores históricos e da fragmentação do habitat sobre a diversidade genética de *Pseudoplatystoma corruscans* (Siluriformes: Pimelodidae). Tese de doutorado. Centro de Ciências Biológicas e da Saúde. Universidade Federal de São Carlos. São Carlos, SP.

Mello, P.H., Venturieri, R.L.L., Honji, R.M. et al., 2009. Threatened fishes of the world: *Pseudoplatystoma corruscans* (Agassiz, 1829) (Siluriformes: Pimelodidae). *Environ Biol Fish* 85, 359–360. <https://doi.org/10.1007/s10641-009-9508-1>

Rodrigues, A.C., N.C.L., dos Santos & L.C., Gomes, 2020. Functionally redundant communities do not show differences in the main environmental drivers of different diversity metrics. *AQUATIC SCIENCES*, v. 82, p. 52. <https://doi.org/10.1007/s00027-020-00727-x>

BAUMGARTNER, M. T., A.G. de Oliveira, A.A., Agostinho, L.C. Gomes, 2018. Fish functional diversity responses following flood pulses in the upper Paraná River floodplain. *ECOLOGY OF FRESHWATER FISH*, v. 27, p. 910-919. <https://doi.org/10.1111/eff.12402>

APPENDIX A - Taxa filtered by sampling methodology in which we applied the rarity index. the result, rr. in crescent order. Spp = species, gri = geographic range index, hsi= habitat specificity index, psi= population size index, rr= rarity index.

spp	gri	hsi	psi	rr
A.affinis	0,985156	0,5	0,00157	0,495575
A.caudima	1	1	1	1
A.crassip	0,290121	0,333333	0,00813	0,210528
A.ellisi	1	1	0,076923	0,692308
A.inermis	0,261433	0,333333	0,011765	0,202177
A.lacustr	0,237675	0,333333	0,000685	0,190565
A.militar	0,217876	0,333333	0,0625	0,20457
A.osteomy	0,201121	0,333333	0,0008	0,178418
A.pantane	0,5092	0,5	0,1	0,369733
A.plagioz	0,86206	1	0,333333	0,731798
A.ucayale	0,343275	0,333333	0,111111	0,262573
Ast.lacus	0,163421	0,333333	0,000729	0,165828
B.gauderi	1	1	1	1
B.hilarii	0,30548	0,5	0,5	0,43516
B.orbigny	0,15381	0,333333	0,009804	0,165649
C.britski	0,145267	0,333333	0,006289	0,16163
C.callich	0,149102	0,333333	0,0625	0,181645
C.gariepi	0,130743	0,333333	0,111111	0,191729
C.haroldo	1	1	0,5	0,833333
C.jaguare	0,803856	0,5	0,021739	0,441865
C.jenynsi	0,124519	0,333333	0,125	0,194284
C.kelberi	0,11886	0,333333	0,002538	0,151577
C.macropo	0,99965	1	0,5	0,833217
C.modestu	0,113693	0,333333	0,022222	0,156416
C.nagelii	0,108957	0,333333	0,017544	0,153278
C.niederl	1	1	0,333333	0,777778
C.paranae	0,113687	0,333333	0,055556	0,167525
C.piquiti	0,100577	0,333333	0,021739	0,151883
Cichla sp	0,137056	0,333333	0,5	0,323463
Cichlaspp	1	1	1	1
Creni spp	1	1	1	1
Cyphocasp	1	1	1	1
E.erythri	0,124531	0,5	0,034483	0,219671
E.triline	0,090174	0,333333	0,004202	0,14257
E.viresce	0,963384	0,5	0,071429	0,511604
F.hahni	1	1	1	1
G.gulo	0,132441	0,5	0,018182	0,216874
G.inaequi	0,087168	0,333333	0,006135	0,142212
G.pantana	0,084357	0,333333	0,25	0,222563
G.parague	0,117566	0,5	0,2	0,272522
G.sveni	0,079245	0,333333	0,001389	0,137989

spp	gri	hsi	psi	rr
G.sylvius	0,076914	0,333333	0,018868	0,143039
Gymnotspp	0,074717	0,333333	0,038462	0,148837
H.aff.her	0,804194	0,5	0,5	0,601398
H.ancistr	0,079199	0,333333	0,018868	0,1438
H.argenti	0,070679	0,333333	0,008333	0,137448
H.cf.albo	0,98898	0,5	0,166667	0,551882
H.cf.iher	0,352383	0,333333	0,05	0,245239
H.cochlio	0,068819	0,333333	0,010309	0,137487
H.commers	0,804194	0,5	0,333333	0,545843
H.hermann	1	1	0,5	0,833333
H.interme	1	1	1	1
H.littora	0,067055	0,333333	0,000825	0,133738
H.mbigua	0,065378	0,333333	0,002375	0,133696
H.microst	1	1	1	1
H.misione	0,085113	0,333333	0,5	0,306149
H.oremacu	0,062265	0,333333	0,006536	0,134045
H.orthono	0,060818	0,333333	0,000828	0,13166
H.platyrh	0,059435	0,333333	0,003226	0,131998
H.regani	0,341972	0,333333	0,003155	0,226153
H.strigat	0,354196	0,333333	0,02	0,235843
H.ternetz	1	1	1	1
H.unitaen	0,058115	0,333333	0,004902	0,132117
Hopli sp2	0,056852	0,333333	0,002183	0,130789
Hopli spp	0,055642	0,333333	0,002188	0,130388
Hypos spp	0,064168	0,333333	0,018519	0,138673
I.labrosu	0,053371	0,333333	0,001698	0,129467
L.araguai	0,95476	0,5	0,5	0,651587
L.frideri	0,052304	0,333333	0,001022	0,128887
L.lacustr	0,051278	0,333333	0,001253	0,128622
L.octofas	0,98898	0,5	0,5	0,662993
L.pectora	0,054951	0,333333	0,009709	0,132664
L.platyme	0,049343	0,333333	0,000304	0,12766
L.prolixa	0,343275	0,333333	0,055556	0,244055
L.rostrat	0,054928	0,333333	0,016129	0,134797
L.striatu	0,98898	0,5	0,142857	0,543946
L.vittatu	1	1	0,111111	0,703704
Leporispp	1	1	1	1
Lorica sp	0,053936	0,333333	0,014286	0,133852
M.aff.int	0,0467	0,333333	0,000437	0,126824
M.bonita	1	1	1	1
M.cf.grac	1	1	1	1
M.foresti	1	1	1	1
M.gra/bon	0,804194	0,5	1	0,768065
M.lippinc	0,045881	0,333333	0,002976	0,127397
M.macroce	0,04509	0,333333	0,025	0,134474

spp	gri	hsi	psi	rr
M.obtusid	0,044326	0,333333	0,012048	0,129902
M.paranan	0,354196	0,333333	0,125	0,270843
M.piavuss	0,043587	0,333333	0,003861	0,126927
M.platanu	0,98898	0,5	0,5	0,662993
M.tiete	0,985156	0,5	0,25	0,578385
O.eigenma	0,047552	0,333333	0,090909	0,157265
O.pintoi	1	1	0,25	0,75
P.amandae	0,048663	0,333333	0,1	0,160665
P.ambrose	0,042181	0,333333	0,000635	0,125383
P.armatul	0,344051	0,333333	0,043478	0,240288
P.avanhan	0,045092	0,333333	0,033333	0,137253
P.cf.falk	0,354196	0,333333	0,019608	0,235712
P.corrusc	0,041512	0,333333	0,005263	0,126703
P.fasciat	0,974978	0,333333	0,011494	0,439935
P.galeatu	0,040863	0,333333	0,000438	0,124878
P.gracili	0,044005	0,333333	0,04	0,139113
P.granulo	0,039625	0,333333	0,002237	0,125065
P.kennedy	0,043051	0,333333	0,011236	0,129207
P.lineatu	0,03846	0,333333	0,00052	0,124104
P.maculat	0,037902	0,333333	0,003436	0,124891
P.manguru	0,98898	0,5	0,2	0,562993
P.mes-hib	0,338376	0,5	0,166667	0,335014
P.mesopot	0,037361	0,333333	0,012346	0,12768
P.microst	0,041845	0,333333	0,25	0,208393
P.mysteri	0,036323	0,333333	0,009615	0,126424
P.nasus	0,98898	0,5	0,1	0,52966
P.ornatus	0,040704	0,333333	0,03125	0,135096
P.paranae	1	1	0,5	0,833333
P.pirinam	0,34376	0,333333	0,01	0,229031
P.reticul	0,95476	0,5	0,071429	0,50873
P.schubar	0,967821	0,5	0,083333	0,517051
P.squamos	0,035341	0,333333	0,002079	0,123585
P.stramin	0,041027	0,5	0,333333	0,291454
P.taeniop	0,804194	0,5	0,333333	0,545843
Pimel spp	0,3393	0,5	0,333333	0,390878
Pimelodsp	1	1	1	1
Pon e vir	0,376598	0,5	0,090909	0,322502
Potamo sp	0,057018	0,5	1	0,519006
Pseudo sp	0,38908	0,5	1	0,629693
R.aspera	0,036324	0,333333	0,00369	0,124449
R.descalv	0,034411	0,333333	0,001008	0,122918
R.dorbign	0,38908	0,5	0,1	0,329693
R.hahni	0,033964	0,333333	0,03125	0,132849
R.quelen	0,309751	0,333333	0,058824	0,233969
R.vulpinu	0,033529	0,333333	0,003906	0,12359

spp	gri	hsi	psi	rr
S.altopar	0,033105	0,333333	0,047619	0,138019
S.borelli	0,032691	0,333333	0,000828	0,122284
S.brasili	0,03353	0,333333	0,006536	0,124467
S.brevipi	0,032287	0,333333	0,000809	0,122143
S.geryi	1	1	1	1
S.hilarii	0,032692	0,333333	0,012987	0,126337
S.insculp	0,031894	0,333333	0,000786	0,122004
S.lima	0,031509	0,333333	0,017241	0,127361
S.macruru	0,034082	0,333333	0,029412	0,132276
S.maculat	0,030768	0,333333	0,002092	0,122064
S.margina	0,03041	0,333333	0,00025	0,121331
S.marmora	1	1	0,5	0,833333
S.nasutus	0,03598	0,333333	0,005556	0,124956
Satanosp	0,030061	0,333333	0,001815	0,121736
Schizo sp	1	1	1	1
Steindasp	1	1	1	1
Steindspp	0,3393	0,5	0,166667	0,335322
T.nematur	1	1	0,0625	0,6875
T.paragua	0,029386	0,333333	0,000904	0,121208
Trache sp	0,02906	0,333333	0,25	0,204131
Z.jahu	0,98898	0,5	0,058824	0,515935

APPENDIX B – Fish species whose functional traits were available.

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>A.affinis</i>	Detritivore	Benthopelagic	Fusiform	Low	93.3	Lateral	Subterminal	Crepuscular	EFNP	Parceled	Sedentary
<i>A.caudima</i>	Insectivore	Benthopelagic	Fus/Lat.compressed	Low	NA	Dorsolateral	Terminal	Nocturnal	EFPC	NA	NA
<i>A.crassip</i>	Piscivore	Benthopelagic	Deep	Extreme	172.3	Lateral	Terminal	Diurnal	EFPC	Parceled	Sedentary
<i>A.ellisi</i>	Insectivore	Benthopelagic	Fus/Lat.compressed	Low	175	Dorsolateral	Terminal	Nocturnal	EFPC	Parceled	Sedentary
<i>A.inermis</i>	Piscivore	Pelagic	Deep	Moderate	212	Lateral	Terminal	Nocturnal	IF	Total	Sedentary
<i>A.lacustr</i>	Piscivore	Benthopelagic	Fusiform	Low	150	Dorsolateral	Terminal	Diu/Noc/Cre	EFNP	Parceled	Sedentary
<i>A.militar</i>	Piscivore	Pelagic	Fusiform	Low	202	Lateral	Terminal	Nocturnal	IF	Total	Sedentary
<i>A.osteomy</i>	Insectivore	Pelagic	Fusiform	Low	191	Lat/Inf	Terminal	Nocturnal	IF	Parceled	Sedentary
<i>A.pantane</i>	Piscivore	Benthopelagic	Fusiform	Low	147	Dorsolateral	Terminal	Diurnal	EFNP	Parceled	Sedentary
<i>A.plagioz</i>	Herbivore	Benthopelagic	Deep/Lat.compressed	Moderate	103	Lateral	Terminal	Diurnal	EFPC	Parceled	NA
<i>A.ucayale</i>	Piscivore	Pelagic	Fusiform	Moderate	290	Lateral	Terminal	Nocturnal	IF	Total	Sedentary
<i>Ast.lacus</i>	Omnivore	Pelagic	Deep	Low	79.9	Lateral	Terminal	Diurnal	EFNP	Parceled	Sedentary
<i>B.gauderi</i>	Insectivore	Benthopelagic	Fus/Lat.compressed	Moderate	NA	Dorsolateral	Terminal	Nocturnal	EFPC	Parceled	Sedentary
<i>B.hilarii</i>	Omnivore	Benthopelagic	Deep	Low	288.9	Lateral	Terminal	Crepuscular	EFNP	Total	Migratory
<i>B.orbigny</i>	Insectivore	Benthopelagic	Fusiform	Low	270	Lateral	Terminal	Crepuscular	EFNP	Total	Migratory
<i>C.britski</i>	Insectivore	Benthopelagic	Deep	Moderate	130	Lateral	Terminal	Diurnal	EFPC	NA	Migratory
<i>C.callich</i>	Insectivore	Demersal	Fusiform	Low	120	Dorsolateral	Terminal	Nocturnal	EFPC	Total	Sedentary
<i>C.gariepi</i>	Piscivore	Benthopelagic	Fusiform	Moderate	27.7	Superior	Terminal	Nocturnal	EFPC	Parceled	Sedentary

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>C.haroldo</i>	Omnivore	Benthopelagic	Fusiform	Moderate	110	Lateral	Terminal	Diurnal	NA	NA	NA
<i>C.jaguare</i>	Omnivore	Benthopelagic	Fusiform	Moderate	101.4	Lateral	Terminal	Diurnal	EFPC	NA	Sedentary
<i>C.jenynsi</i>	Insectivore	Demersal	Deep/Laterally compressed	Extreme	107.6	Lateral	Terminal	Nocturnal	EFNP	Parceled	NA
<i>C.kelberi</i>	Piscivore	Benthopelagic	Deep	Moderate	190	Dorsolateral	Terminal	Diurnal	EFPC	Total	Sedentary
<i>C.macropo</i>	Omnivore	Benthopelagic	Deep	Low	250	Lateral	Terminal	Diurnal	EFNP	Total	Migratory
<i>C.modestu</i>	Detritivore	Benthopelagic	Deep	Low	80	Lateral	Terminal	Diurnal/Noc	EFNP	Total	Sedentary
<i>C.nagelii</i>	Detritivore	Benthopelagic	Fusiform	Low	130	Lateral	Terminal	Diurnal/Noc/Cree	EFNP	Total	Sedentary
<i>C.niederl</i>	Insectivore	Benthopelagic	Fusiform	Low	145.5	Lateral	Terminal	Diurnal	EFPC	NA	Sedentary
<i>C.paranae</i>	Piscivore	Benthopelagic	Deep	Moderate	110	Dorsolateral	Terminal	Diurnal	EFPC	Parceled	Sedentary
<i>C.piquiti</i>	Piscivore	Benthopelagic	Deep	Moderate	240	Dorsolateral	Terminal	Diurnal	EFPC	Parceled	Sedentary
<i>E.erythri</i>	Piscivore	Demersal	Fusiform/Cylindrical	Low	93	Dorsolateral	Terminal	Diurnal	EFPC	Total	Sedentary
<i>E.triline</i>	Insectivore	Benthopelagic	Fus/Dee	Moderate	212	Dorsolateral	Terminal	Crepuscular	EFPC	Total	NA
<i>E.viresce</i>	Insectivore	Benthopelagic	Fusiform	Extreme	275	Dorsolateral	Subterminal	Crepuscular	EFNP	Parceled	Sedentary
<i>F.hahni</i>	Planktivore	Demersal	Flattened	Low	130	Dorsolateral	Inferior	Cre/Noc	EFPC	NA	NA
<i>G.gulo</i>	Piscivore	Benthopelagic	Deep	Low	200	Lateral	Sub/Ter	Crepuscular	EFNP	Continuou s	Sedentary
<i>G.inaequi</i>	Insectivore	Benthopelagic	Fusiform	Moderate	134	Ventrolateral	Superior	Nocturnal	EFPC	Parceled	Sedentary
<i>G.pantana</i>	Omnivore	Benthopelagic	Fusiform	Moderate	NA	Ventrolateral	Superior	Nocturnal	IF	Total	Sedentary
<i>G.parague</i>	Omnivore	Benthopelagic	Fusiform	Low	240	Ventrolateral	Superior	Crepuscular	IF/EFPC	Total	Sedentary
<i>G.sveni</i>	Invertivore	Demersal	Deep	Low	150	Lateral	Terminal	Diurnal	EFPC	NA	Sedentary
<i>G.sylvius</i>	Invertivore	Benthopelagic	Fusiform	Low	175.4	Ventrolateral	Superior	Nocturnal	EFPC	Total	Sedentary

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>H.aff.her</i>	Detritivore	Demersal	Deep	Moderate	NA	Dorsolateral	Inferior	Crepuscular	EFPC	Total	Sedentary
<i>H.ancistr</i>	Detritivore	Demersal	Deep	Extreme	140	Dorsolateral	Inferior	Noc/Cre	EFPC	Total	Sedentary
<i>H.cf.albo</i>	Detritivore	Demersal	Deep	Moderate	210	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.cochlio</i>	Detritivore	Demersal	Deep	Moderate	230	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.commers</i>	Detritivore	Demersal	Deep	Low	255	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.interme</i>	Piscivore	Pelagic	Fus/Cyl	Moderate	277.2	Lateral	Terminal	Cre/Noc	EFPC	Parceled	Sedentary
<i>H.littora</i>	Invertivore	Demersal	Fusiform	Low	133.4	Lateral	Terminal	Cre/Noc	EFPC	Parceled	Sedentary
<i>H.mbigua</i>	Piscivore	Pelagic	Fus/Cyl	Moderate	260	Lateral	Terminal	Cre/Noc	EFPC	Parceled	Sedentary
<i>H.microst</i>	Detritivore	Demersal	Deep	Moderate	165	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.oremacu</i>	Invertivore	Pelagic	Fusiform	Moderate	405	Ventrolateral	Terminal	Diu/Noc	EFNP	Parceled	Sedentary
<i>H.orthono</i>	Omnivore	Pelagic	Fusiform	Low	137	Lateral	Terminal	Diu/Cre	EFNP	Total	Sedentary
<i>H.platyrrh</i>	Piscivore	Demersal	Fusiform	Moderate	470	Superior	Terminal	Cre/Noc	EFNP	Total	Migratory
<i>H.regani</i>	Detritivore	Demersal	Deep	Moderate	170	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.strigat</i>	Detritivore	Demersal	Deep	Moderate	150	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.ternetz</i>	Detritivore	Demersal	Deep	Low	210	Dorsolateral	Inferior	Cre/Noc	EFPC	Total	Sedentary
<i>H.unitaen</i>	Omnivore	Benthopelagic	Fus/Cyl	Moderate	150.2	Dorsolateral	Terminal	Diu/Noc	EFPC	NA	Sedentary
<i>Hopli.sp2</i>	Piscivore	Pelagic	Fus/Cyl	Low	250	Lateral	Terminal	Cre/Noc	EFPC	Parceled	Sedentary
<i>I.labrosu</i>	Invertivore	Benthopelagic	Fusiform	Moderate	200	Dorsolateral	Subterminal	Cre/Noc	EFNP	Parceled	Sedentary
<i>L.araguai</i>	Insectivore	Benthopelagic	Deep	Moderate	32.6	Lateral	Terminal	Diurnal	EFPC	NA	NA

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>L.frideri</i>	Omnivore	Benthopelagic	Fusiform	Low	189.2	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Sedentary
<i>L.lacustr</i>	Omnivore	Benthopelagic	Deep	Low	210	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Parceled	Sedentary
<i>L.octofas</i>	Omnivore	Benthopelagic	Deep	Moderate	148.9	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Sedentary
<i>L.pectora</i>	Invertivore	Benthopelagic	Fusiform	Moderate	100	Dorsolateral	Terminal	Nocturnal	EFPC	NA	Sedentary
<i>L.platyme</i>	Detritivore	Demersal	Flattened	Low	210	Superior	Inferior	Diu/Noc/Cr e	EFPC	Parceled	Sedentary
<i>L.prolixa</i>	Herbivore	Demersal	Flattened	Moderate/Extreme	230.5	Dorsolateral	Inferior	Diu/Noc/Cr e	EFPC	NA	Sedentary
<i>L.rostrat</i>	Detritivore	Demersal	Flattened	Low	250	Superior	Inferior	Diu/Noc/Cr e	EFPC	Parceled	Sedentary
<i>L.striatu</i>	Insectivore	Benthopelagic	Fusiform	Moderate	69.2	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	NA
<i>L.vittatu</i>	Insectivore	Benthopelagic	Fusiform	Low	200	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Sedentary
<i>M.aff.int</i>	Invertivore	Pelagic	Fusiform	Low	68.4	Lateral	Terminal	Diu/Noc	EFNP	Parceled	Sedentary
<i>M.bonita</i>	Insectivore	Pelagic	Lat. compressed	Low	28.-.44	Lateral	Terminal	Diu/Noc	EFNP	Parceled	Sedentary
<i>M.cf.grac</i>	Insectivore	Pelagic	Deep	Low	39.7	Lateral	Terminal	Diu/Noc	EFNP	Parceled	Sedentary
<i>M.foresti</i>	Herbivore	Pelagic	Lat. compressed	Low	30.6	Lateral	Terminal	Diu/Noc	EFNP	Parceled	Sedentary
<i>M.lippinc</i>	Herbivore	Pelagic	Deep/Lat.compressed	Low	149.3	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Sedentary
<i>M.macroce</i>	Omnivore	Benthopelagic	Deep	Moderate	470	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Migratory
<i>M.obtusid</i>	Omnivore	Benthopelagic	Fusiform	Low	190.8	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Migratory
<i>M.paranan</i>	Detritivore	Demersal	Deep	Low	205	Dorsolateral	Inferior	Diu/Noc/Cr e	EFPC	Total	Sedentary
<i>M.plavuss</i>	Omnivore	Benthopelagic	Fusiform	Low	190.8	Lateral	Terminal	Diu/Noc/Cr e	EFNP	Total	Migratory
<i>M.platanu</i>	Piscivore	Demersal	Fusiform	Low	340	Dorsolateral	Terminal	Nocturnal	NA	NA	NA

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>M.tiete</i>	Herbivore	Benthopelagic	Deep/Lat.compressed	Low	135	Lateral	Terminal	Diu/Noc/Cre	EFNP	NA	Sedentary
<i>O.eigenma</i>	Insectivore	Demersal	Deep	Extreme	82.9	Dorsolateral	Terminal	Cre/Noc	NA	NA	NA
<i>O.pintoi</i>	Invertivore	Pelagic	Deep	Extreme	68.4	Lateral	Terminal	Cre/Noc	EFNP	Parceled	Sedentary
<i>P.amandae</i>	Piscivore	Demersal	Flattened	Extreme	380	Superior	Inferior	Diu/Noc	IF	NA	Sedentary
<i>P.ambrose</i>	Detritivore	Demersal	Deep	Low	395	Dorsolateral	Inferior	Crepuscular	EFPC	Parceled	Sedentary
<i>P.armatulus</i>	Invertivore	Demersal	Deep	Low	132.9	Dorsolateral	Terminal	Nocturnal	EFNP	NA	Sedentary
<i>P.avanhan</i>	Invertivore	Demersal	Fusiform	Low	125	Dorsolateral	Terminal	Cre/Noc	EFNP	Total	Sedentary
<i>P.cf.falk</i>	Piscivore	Demersal	Flattened	Extreme	NA	Superior	Inferior	Diu/Noc	IF	NA	Sedentary
<i>P.corrusc</i>	Piscivore	Demersal	Fusiform	Low	675	Superior	Subterminal	Cre/Noc	EFNP	Total	Migratory
<i>P.galeatu</i>	Omnivore	Pelagic	Deep	Extreme	165	Dorsolateral	Terminal	Nocturnal	IF	Parceled	Sedentary
<i>P.gracili</i>	Insectivore	Demersal	Fusiform	Moderate	120	Dorsolateral	Terminal	Cre/Noc	EFNP	Total	Sedentary
<i>P.granulo</i>	Omnivore	Demersal	Deep	Extreme	600	Superior	Terminal	Nocturnal	EFNP	Parceled	Migratory
<i>P.kennedy</i>	Herbivore	Benthopelagic	Deep/Lat.compressed	Moderate	30.5	Lateral	Terminal	Diurnal	NA	NA	NA
<i>P.lineatu</i>	Detritivore	Benthopelagic	Deep/Lat.compressed	Moderate	495.2	Lateral	Terminal	Diu/Noc/Cre	EFNP	Total	Migratory
<i>P.maculatus</i>	Omnivore	Demersal	Deep	Moderate	180	Dorsolateral	Terminal	Crepuscular	EFNP	Parceled	Migratory
<i>P.manguru</i>	Detritivore	Demersal	Fusiform/Deep	Moderate	180	Superior	Terminal	Crepuscular	NA	NA	NA
<i>P.mesopot</i>	Herbivore	Demersal	Deep/Lat.compressed	Low	498	Lateral	Terminal	Diu/Noc/Cre	EFNP	Total	Migratory
<i>P.microst</i>	Omnivore	Demersal	Deep	Moderate	140	Dorsolateral	Terminal	Crepuscular	EFNP	Total	Sedentary
<i>P.mysteri</i>	Omnivore	Demersal	Deep	Low	90.4	Dorsolateral	Terminal	Crepuscular	EFNP	Total	NA
<i>P.nasus</i>	Planktivore	Benthopelagic	Fusiform	Moderate	102.5	Lateral	Subterminal	Cre/Noc	EFNP	Parceled	Sedentary
<i>P.ornatus</i>	Omnivore	Demersal	Deep	Low	208	Dorsolateral	Terminal	Crepuscular	EFNP	Total	Migratory

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>P.paranae</i>	Insectivore	Pelagic	Fus/Lat.compressed	Moderate	NA	Lateral	Terminal	Diurnal	EFNP	NA	Sedentary
<i>P.pirnam</i>	Piscivore	Demersal	Deep	Extreme	620	Dorsolateral	Terminal	Crepuscular	EFNP	Total	Migratory
<i>P.reticul</i>	Piscivore	Demersal	Fusiform	Moderate	580	Superior	Subterminal	Cre/Noc	EFNP	Total	Migratory
<i>P.schubar</i>	Insectivore	Pelagic	Fus/Lat.compressed	Moderate	NA	Lateral	Terminal	Diurnal	EFNP	NA	Sedentary
<i>P.squamos</i>	Piscivore	Pelagic	Deep	Low	330	Dorsolateral	Terminal	Nocturnal	EFNP	Parceled	Sedentary
<i>P.stramin</i>	Insectivore	Pelagic	Fusiform	Low	70	Lateral	Terminal	Crepuscular	EFNP	Total	Sedentary
<i>P.taeniop</i>	Insectivore	Demersal	Fusiform	Low	130	Dorsolateral	Terminal	Cre/Noc	EFNP	Total	Sedentary
<i>R.aspera</i>	Detritivore	Demersal	Deep	Low	360	Dorsolateral	Inferior	Crepuscular	EFNP	Total	Migratory
<i>R.descalv</i>	Insectivore	Benthopelagic	Deep	Low	88.8	Lateral	Terminal	Cre/Noc	EFNP	Parceled	Sedentary
<i>R.dorbign</i>	Invertivore	Demersal	Deep	Low	135.5	Dorsolateral	Subterminal	Nocturnal	EFNP	NA	Sedentary
<i>R.hahni</i>	Insectivore	Benthopelagic	Fus/Lat.compressed	Moderate	700	Dorsolateral	Terminal	Nocturnal	EFPC	Parceled	Sedentary
<i>R.quelen</i>	Piscivore	Demersal	Fusiform	Extreme	163.2	Dorsolateral	Terminal	Cre/Noc	EFNP	Parceled	Sedentary
<i>R.vulpinu</i>	Piscivore	Pelagic	Fusiform	Low	480	Dorsolateral	Superior	Cre/Noc	EFNP	Total	Migratory
<i>S.altopar</i>	Herbivore	Benthopelagic	Fusiform	Low	282	Lateral	Terminal	Diu/Noc/Cre	EFNP	Total	Sedentary
<i>S.borelli</i>	Herbivore	Benthopelagic	Fusiform	Low	207.5	Lateral	Terminal	Diu/Noc/Cre	EFNP	Total	Sedentary
<i>S.brasili</i>	Piscivore	Benthopelagic	Deep	Moderate	500	Lateral	Terminal	Diurnal	EFNP	Total	Migratory
<i>S.brevipi</i>	Detritivore	Benthopelagic	Fusiform	Moderate	105.8	Lateral	Terminal	Diu/Noc/Cre	EFNP	NA	Sedentary
<i>S.hilarii</i>	Piscivore	Benthopelagic	Deep	Moderate	170	Lateral	Terminal	Diurnal	EFNP	Total	Migratory
<i>S.insculp</i>	Detritivore	Pelagic	Fusiform	Moderate	90	Lateral	Terminal	Diu/Noc/Cre	EFNP	Total	Sedentary
<i>S.lima</i>	Piscivore	Demersal	Fusiform	Low	487	Lateral	Subterminal	Cre/Noc	EFNP	Total	Migratory

spp	Trophic category	Position in the water column	Body shape	Crypsis	Standard length (mm)	Eye position	Mouth position	Diel activity	Fecundation and parental care	Spawning	Migration
<i>S.macruru</i>	Insectivore	Benthopelagic	Fusiform	Moderate	625	Dorsolateral	Terminal	Crepuscular	EFPC/EFNP	Parceled	Sedentary
<i>S.maculat</i>	Piscivore	Pelagic	Deep/Lat.compressed	Moderate	157.2	Lateral	Terminal	Diu/Noc/Cre	EFPC	Parceled	Sedentary
<i>S.margina</i>	Piscivore	Pelagic	Deep/Lat.compressed	Moderate	160.2	Lateral	Terminal	Diu/Noc/Cre	EFPC	Parceled	Sedentary
<i>S.marmora</i>	Herbivore	Benthopelagic	Anguiliform	Moderate	273.2	Lateral	Terminal	Cre/Noc	EFPC	NA	Sedentary
<i>S.nasutus</i>	Herbivore	Pelagic	Fusiform	Low	160	Lateral	Subterminal	Diu/Noc/Cre	EFNP	Total	Sed/Mig
<i>Satanop.sp</i>	Detritivore	Benthopelagic	Deep	Extreme	150	Dorsolateral	Terminal	Diurnal	EFPC	Parceled	Sedentary
<i>T.nematur</i>	Insectivore	Benthopelagic	Deep	Low	102	Lateral	Terminal	Diurnal	EFNP	NA	Sedentary
<i>T.paragua</i>	Invertivore	Demersal	Deep	Moderate	125	Dorsolateral	Subterminal	Nocturnal	EFNP	Parceled	Sedentary
<i>Trache.sp</i>	Omnivore	Pelagic	Deep	Moderate	98.2	Dorsolateral	Terminal	Nocturnal	IF	NA	Sedentary
<i>Z.jahu</i>	Piscivore	Demersal	Deep	Moderate	800	Dorsal	Terminal	Cre/Noc	EFNP	Total	Migratory

APPENDIX C – Functional rarity index results.

spp	Uniqueness	Restrictedness	Functional rarity
M.obtusid	0.00	0	0
M.piavuss	0	0	0
R.hahni	0	0	0
C.piquiti	0.09090909	0	0.045454545
H.mbigua	0.09090909	0	0.045454545
L.platyme	0.09090909	0	0.045454545
S.altopar	0.09090909	0	0.045454545
S.borelli	0.09090909	0	0.045454545
S.maculat	0.09090909	0	0.045454545
S.margina	0.09090909	0	0.045454545
P.mysteri	0.1	0	0.05
C.britski	0.11111111	0	0.055555555
A.lacustr	0.18181818	0	0.09090909
C.kelberi	0.18181818	0	0.09090909
Hopli.sp2	0.18181818	0	0.09090909
L.frideri	0.18181818	0	0.09090909
M.macroce	0.18181818	0	0.09090909
P.corrusc	0.18181818	0	0.09090909
S.lima	0.18181818	0	0.09090909
M.lippinc	0.2	0	0.1
P.ambrose	0.2	0	0.1
P.galeatu	0.2	0	0.1
R.descalv	0.2	0	0.1
S.brevipi	0.2	0	0.1
S.insculp	0.2	0	0.1
T.paragua	0.2	0	0.1
H.cochlio	0.09090909	0.11111111	0.101010095
L.rostrat	0.09090909	0.11111111	0.101010095
S.brasili	0.09090909	0.11111111	0.101010095
A.crassip	0.27272727	0	0.136363635
Ast.lacus	0.27272727	0	0.136363635
B.orbigny	0.27272727	0	0.136363635
H.oremacu	0.27272727	0	0.136363635
H.orthono	0.27272727	0	0.136363635
L.lacustr	0.27272727	0	0.136363635

spp	Uniqueness	Restrictedness	Functional rarity
M.aff.int	0.27272727	0	0.136363635
P.lineatu	0.27272727	0	0.136363635
P.maculat	0.27272727	0	0.136363635
P.mesopot	0.27272727	0	0.136363635
Satanop.sp	0.27272727	0	0.136363635
A.militar	0.18181818	0.11111111	0.14646464
H.ancistr	0.18181818	0.11111111	0.14646464
H.platyrh	0.18181818	0.11111111	0.14646464
G.inaequi	0.3	0	0.15
G.sveni	0.3	0	0.15
H.littora	0.3	0	0.15
C.paranae	0.09090909	0.22222222	0.156565645
P.amandae	0	0.33333333	0.16666665
L.pectora	0.22222222	0.11111111	0.16666666
E.triline	0.33333333	0	0.166666665
H.unitaen	0.33333333	0	0.166666665
A.osteomy	0.36363636	0	0.18181818
R.vulpinu	0.36363636	0	0.18181818
C.modestu	0.27272727	0.11111111	0.191919185
A.inermis	0.18181818	0.22222222	0.20202019
P.avanhan	0.18181818	0.22222222	0.20202019
P.gracili	0.18181818	0.22222222	0.20202019
G.sylvius	0.3	0.11111111	0.20555555
P.squamos	0.3	0.11111111	0.20555555
H.regani	0.09090909	0.33333333	0.212121195
P.ornatus	0.1	0.33333333	0.21666665
C.callich	0.25	0.22222222	0.23611111
I.labrosu	0.36363636	0.11111111	0.23737373
P.granulo	0.36363636	0.11111111	0.23737373
R.aspera	0.27272727	0.22222222	0.247474735
C.nagelii	0.18181818	0.33333333	0.25757574
C.gariepi	0.3	0.22222222	0.26111111
S.macruru	0.3	0.22222222	0.26111111
Trache.sp	0.2	0.33333333	0.26666665
S.hilarii	0.09090909	0.44444444	0.267676745
P.kennedy	0.125	0.44444444	0.2847222
O.eigenma	0.375	0.22222222	0.2986111

spp	Uniqueness	Restrictedness	Functional rarity
C.jenynsi	0.4	0.2222222	0.3111111
A.pantane	0.18181818	0.4444444	0.31313129
A.ucayale	0.18181818	0.4444444	0.31313129
G.pantana	0.3	0.3333333	0.31666665
P.microst	0.2	0.4444444	0.3222222
P.cf.falk	0	0.6666667	0.33333335
P.schubar	0	0.6666667	0.33333335
G.parague	0.3	0.4444444	0.3722222
P.armatul	0.2	0.5555556	0.3777778
H.strigat	0.09090909	0.6666667	0.378787895
L.araguai	0.11111111	0.6666667	0.388888905
C.jaguare	0.125	0.6666667	0.39583335
E.erythri	0.36363636	0.4444444	0.40404038
E.viresce	0.36363636	0.4444444	0.40404038
R.quelen	0.36363636	0.4444444	0.40404038
S.nasutus	0.36363636	0.4444444	0.40404038
P.pirinam	0.27272727	0.5555556	0.414141435
G.gulo	0.4	0.4444444	0.4222222
L.prolixa	0.4	0.4444444	0.4222222
M.paranan	0.18181818	0.6666667	0.42424244
M.tiete	0.2	0.6666667	0.43333335
H.cf.albo	0.09090909	0.7777778	0.434343445
H.commers	0.09090909	0.7777778	0.434343445
H.aff.her	0.1	0.7777778	0.4388889
A.caudima	0	0.8888889	0.44444445
A.ellisi	0	0.8888889	0.44444445
B.gauderi	0	0.8888889	0.44444445
P.paranae	0	0.8888889	0.44444445
A.plagioz	0.125	0.7777778	0.4513889
P.stramin	0.27272727	0.6666667	0.469696985
B.hilarii	0.18181818	0.7777778	0.47979799
C.macropo	0.18181818	0.7777778	0.47979799
L.octofas	0.18181818	0.7777778	0.47979799
P.reticul	0.18181818	0.7777778	0.47979799
P.taeniop	0.18181818	0.7777778	0.47979799
L.striatu	0.2	0.7777778	0.4888889
R.dorbign	0.2	0.7777778	0.4888889

spp	Uniqueness	Restrictedness	Functional rarity
H.interme	0.09090909	0.8888889	0.489898995
H.microst	0.09090909	0.8888889	0.489898995
H.ternetz	0.09090909	0.8888889	0.489898995
C.haroldo	0.125	0.8888889	0.50694445
M.platanu	0.25	0.7777778	0.5138889
A.affinis	0.36363636	0.67	0.51515153
Z.jahu	0.27272727	0.7777778	0.525252535
L.vittatu	0.18181818	0.8888889	0.53535354
M.bonita	0.18181818	0.8888889	0.53535354
M.cf.grac	0.18181818	0.8888889	0.53535354
M.foresti	0.18181818	0.8888889	0.53535354
T.nematur	0.2	0.8888889	0.54444445
P.nasus	0.36363636	0.7777778	0.57070708
P.manguru	0.375	0.7777778	0.5763889
O.pintoi	0.27272727	0.8888889	0.580808085
C.niederl	0.3	0.8888889	0.59444445
S.marmora	0.33	0.8888889	0.611111115
F.hahni	0.44444444	0.8888889	0.66666667